

Original Research

Assessment of Lead and Cadmium Bioaccumulation by Tilapia, *Sarotherodon melanotheron* (Rüppell, 1852), from the Lake Nokoué Lagoon of Porto-Novo Complex, Benin Republic

Gisèle Houédjissi^{1,2,3}, Maxime Machioud Sangaré-oumar^{1,2†}, Prudencio Agbohessi³, Chabi Oloushegun Onibon³ and Jean-Michel Akpo¹

¹ Laboratory of Pharmacology and Experimental Physiology (LPPE), Faculty of Science and Technology, University of Abomey-Calavi, Benin

² National Water Institute (INE), African Center of Excellence for Water and Sanitation, University of Abomey-Calavi, Benin

³ Research Laboratory in Aquaculture and Aquatic Ecotoxicology (LaRAEq), University of Parakou, Research Laboratory in Aquaculture and Aquatic Ecotoxicology, Benin

† Corresponding author: Maxime Machioud Sangaré-Oumar; sangoumarfr@yahoo.fr

Gisèle Houédjissi: ORCID ID <https://orcid.org/0009-0005-3542-3796>

Maxime Machioud Sangaré-Oumar: ORCID ID <https://orcid.org/0000-0001-7081-4120>

Prudencio Agbohessi: ORCID ID <https://orcid.org/0000-0002-3922-2167>

Key Words	Lead, cadmium, bioaccumulation, Lake Nokoué Porto-Novo lagoon complex, <i>Sarotherodon melanotheron</i>
DOI	https://doi.org/10.46488/NEPT.2025.v24i04.D1806 (DOI will be active only after the final publication of the paper)
Citation for the Paper	Houédjissi, G., Sangaré-oumar, M.M., Agbohessi, P., Onibon, C.O. and Akpo, J. M., 2025. Assessment of lead and cadmium bioaccumulation by Tilapia, <i>Sarotherodon melanotheron</i> (Rüppell, 1852), from the Lake Nokoué lagoon of Porto-Novo complex, Benin Republic. <i>Nature Environment and Pollution Technology</i> , 24(4), D1806. https://doi.org/10.46488/NEPT.2025.v24i04.D1806

ABSTRACT

Lake Nokoué-lagoon of Porto-Novo complex, as a surface water body, is exposed to heavy pollution whose intensity vary across seasons. The present study aimed to assess the bioaccumulation of lead and cadmium levels by *Sarotherodon melanotheron* fish, one of the most populated fished in this complex. To this end, fish (muscle, gills, female gonads), water and sediment were sampled during high-water and low-water. Lead and cadmium were quan-

tified from different samples using a flame atomic absorption spectrometry. Health risk associated to the fish consumption was estimated. In the water samples, lead concentration was higher during high water (0.106 ± 0.116 mg/l); cadmium was higher during low water (0.010 ± 0.001 mg/l). In the sediment samples, lead was higher during high water (13.94 ± 20.79 mg/kg); cadmium was higher during low water (0.2 ± 0.06 mg/kg). On the other hand, high concentrations of lead and cadmium in fish were found during low water, respectively 27.66 mg/kg and 0.22 mg/kg in the muscle. cadmium accumulation in fish is influenced by dissolved oxygen and the pH of the water. Consumption of 400 g of *Sarotherodon melanotheron* fish from this complex per week constitutes a health risk for anyone weighing between 0 and 90 kg.

INTRODUCTION

The Lake Nokoué lagoon of Porto-Novo complex, later referred to as “the Complex” is one of the most important fishing areas in West Africa (Lalèyè *et al.*, 1995) and accounts for more than half of continental fish production in Benin (65-70%) (Lalèyè *et al.*, 2004). However, this Complex is surrounded by Benin's most densely populated municipalities, with around 2 million inhabitants living around it (INSAE, 2015), and the majority of Benin's industries are located near the Complex (Elegbede Manou *et al.*, 2020). Therefore, the Complex constitutes a receptacle for industrial discharges, which are the major source of heavy metals and all the solid and liquid wastes of neighboring populations. The Complex is also the main route for petroleum products, particularly leaded petrol, from Nigeria to Benin (Dovonou, 2019). All this exposes the Complex to heavy pollution from human activities. Various wastes generated by the surrounding population, including domestic, hospital and industrial wastes, and petroleum products, are the main sources of heavy metals (Rossi, 2009). Heavy metals with a density greater than 5 g/cm³, such as cadmium and lead, are classified as toxic (Cuniasse and Glass, 2020). The presence of these toxic metals in an ecosystem exposes it to a high risk of degradation. Studies have shown the capacity for bioaccumulation and biomagnification of these toxic metals in aquatic fauna, particularly fish, due to their lipophilic nature, and top links are the most vulnerable (Başyigit and Tekin-özan, 2013). Most of these heavy metals are also endocrine disruptors for fish (Barbier, 2011) and also have carcinogenic effects on human health. Despite previous studies that focused on this Complex (Agbandou *et al.*, 2018; Elegbede Manou *et al.*, 2020; Agbohessi *et al.*, 2023a), the seasonal pollution level, the bioaccumulation across various organs in the fish and the health risk assessment relatively to the consumption of *Sarotherodon melanotheron* from the Complex remain largely unknown. Therefore, this study was undertaken to fill this gap of knowledge in order to stimulate the public and decision-makers awareness on the destruction of biodiversity and the health risk for the population.

2. MATERIALS AND METHODS

The study was carried out during the high-water and low water seasons, in March 2023 and October 2024. To this end, sampling sites were selected, samples were collected, lead and cadmium level were determined, and the data were statically processed.

2.1 Sampling sites

The complex is located in south-east Benin (West Africa) and covers an area of 180 km² at low water (Figure 1). Lake Nokoué (160 km² at low water) and the Porto-Novo lagoon (20 km² at low water) are a brackish water complex located between parallels 6°25' and 6°30' North and meridians 2°20' and 2°40' East, linked by the Totché channel (Colleuil & Texier, 1987). Fed by freshwater from the Ouémé river, the Sô river, rainfall and run-off, Lake Nokoué is connected to the Atlantic Ocean by the Cotonou channel over a length of 4.5 km and a width of around 300 m (Lalèye et al., 1995).

Fifteen (15) sites were identified on the complex for sampling (Figure 1). These sites were identified for the following reasons:

Proximity to housing

Proximity to human activities (markets, leisure facilities, fishing areas, etc.)

Proximity to hospitals




Figure 1: Representation of sampling sites

2.2 Samples collection

During sampling, number of physico-chemical parameters, such as temperature, dissolved oxygen, pH and salinity, were measured at each site in order to provide a general characteristic of the Complex. These physico-chemical parameters were taken in the morning from 7am to 9am and in the evening from 4 pm to 6pm at all sites to a depth of 30 cm. Water samples were taken in 1-litre bottles and sediment samples were taken in jars using a sampling bucket at each of the identified sites. Specimens of *Sarotherodon melanotheron* at various sites to collect muscle, gonads and gills for metal assays. Eighty (80) fish were collected during each season, making a total of 160 fish. Samples are stored in an isothermal cooler containing ice in the field. In the laboratory, water samples were acidified with nitric acid. Sediment, female gonads, gills and muscle were oven-dried at 99°C for lead and cadmium determination.

2.3 Lead and cadmium analysis

Sample preparation: lead and cadmium in water were determined using the atomic absorption method described above (Rodier et al., 2009). Water samples were filtered at 0.45 µm, acidified to 1% with HNO₃ and stored at room temperature until analysis.

For sediment, the acid-etch mineralization method (a mixture of hydrochloric acid and nitric acid) was used. One gram (1g) of sediment was mixed with 3mL of 37% hydrochloric acid and 1mL of 65% nitric acid at 110-150 °C. The mixture is then filtered to obtain the filtrate which will be used for the determination of these metals.

For fish organ samples, 1g of organ was immersed in 10mL of 9% hydrogen peroxide (H₂O₂) for 24 hours. Then 4mL of 65% nitric acid was added, heated to 150°C for 2 minutes and filtered to obtain the filtrate.

Determination of concentration: A flame atomic absorption spectrometer (SAA Ice 3000 SERIES THERMO FISCHER) was used for the determination of lead and cadmium in the previously prepared filtrates. After establishing the calibration curve from the standards, the “optical densities” were read to obtain the concentrations of the elements in the filtrates, then related to the mass of sediment or fish. These concentrations were determined under the following experimental conditions (Table 1). The spectrometer displays the concentration of the element studied in the samples directly to the computer. These concentrations are expressed in mg/l. However, the actual concentrations (C) of the elements measured in solid samples are expressed in mg/Kg of dry weight.

Table 1: Summary of experimental conditions for the analysis of each element

Element analyzed	Wavelength (nm)	Width of slot	Type of flame
Cadmium	228,8	0,5	Air-Acetylene
Lead	217,0	0,5	Air-Acetylene

2.4 Bioaccumulation factor

An estimate of the bioaccumulation factor (Bf) of toxic metals was made in order to see at what rate these metals are accumulated in fish in proportion to the water concentration. The formula used is as follows:

$$Bf = CMF / CMW$$

Bf being the bioaccumulation factor, CMF the concentration of metals in fish in mg/kg and CMW the concentration of metals in water in mg/l.

2.5 Assessment of the health risk associated with fish consumption

According to information gathered in the study area, a local resident may consume two to three *Sarotherodon melanotheron* fish per week estimated at around 400g per week per person. This information was used to calculate Weekly Exposure Doses (WED) for lead and cadmium using the method used by (Elegbede Manou *et al.*, 2020) compared with the Maximum Tolerable Weekly Intake (MTHI) according to the World Health Organisation.

$$WED = (Q * C) / W$$

Q being the Quantity of fish consumed per week by one inhabitant in Kg, C the concentration of lead or cadmium in one Kg in the fish sample in mg/kg and W the weight of one inhabitant in Kg.

2.6 Statistical analysis

The data was entered into the Excel 2016 spreadsheet and processed using XLstat statistical software. The experimental units are water, sediment and fish, and the factor is the high and low-water seasons. The Ryan-Joiner test and the Levene test were used to assess the normality and homoscedasticity of the data, respectively. Data that does not follow a normal distribution is transformed into $\log(x+1)$. Results are expressed as mean \pm standard deviation of the mean. The differences between the means were evaluated using a test based on one-factor analysis of variance (ANOVA) according to the effect studied. Average values were compared between seasons. Duncan's test was used to show a significant difference between the data. A correlation matrix was drawn up between the variables used. For the analyses, a difference was considered statistically significant if the p-value was less than 0.05.

3. RESULTS

3.1. Physico-chemical parameters

The average values of the physico-chemical parameters of the Lake Nokoué-Porto-Novo lagoon complex are presented in Table 2. The average temperature was higher during low water (33.41 and 32.87°C) in the Lake Nokoué-

Porto-Novo lagoon complex than during high water. The same observations were made for dissolved oxygen, pH and salinity. Dissolved oxygen recorded in the Porto-Novo lagoon during both seasons was also low (2.11 and 2.18 respectively during flood and low water).

Table 2: Physico-chemical parameters

Physico-chemical parameters	Seasons	Lake Nokoué	Porto-Novo Lagoon
Water temperature (°C)	High water	29,02 ± 6,27	30,13 ± 0,48
	Low water	33,41 ± 0,74	32,87 ± 0,15
Dissolved oxygen (mg/l)	High water	3,74 ± 6,84	2,11 ± 1,40
	Low water	5,38 ± 1,45	2,18 ± 1,86
pH	High water	6,52 ± 0,24	6,21 ± 0,12
	Low water	8,63 ± 0,54	7,42 ± 0,49
Salinity (g/l)	High water	0,49 ± 0,70	0,68 ± 0,49
	Low water	19,64 ± 10,77	4,83 ± 0,71

3.2 Lead and cadmium concentrations in water and sediment

The concentration of lead and cadmium in water and sediment was expressed as the mean plus or minus standard deviation (mean ± SD, Figures 3 and 4). The high concentration of lead recorded in the water during the flood was 0.106 ± 0.116 mg/l, and that in the sediment was 13.94 ± 20.79 mg/kg during the flood. The high concentration of cadmium in the water was 0.017 ± 0.013 mg/l during the high water event, and that in the sediment was 0.20 ± 0.12 mg/kg during the high water event. The highest concentrations of lead were recorded in the Porto-Novo lagoon and cadmium in Lake Nokoué during the high water.






Figure 2: Lead and cadmium concentrations in water**Figure 3:** Concentration of lead and cadmium in sediment

3.3 Lead and cadmium concentrations in *Sarotherodon melanotheron* fish

The mean concentration of lead and cadmium in the muscle, gills and female gonads of *Sarotherodon melanotheron* is shown in Figure 5. High lead concentrations in muscle, gills and female gonads are 16.41 ± 15.91 mg/kg, 7.93 ± 3.43 mg/kg and 5.277 ± 1.30 mg/kg respectively. Cadmium levels in female muscle, gills and gonads are 0.148 ± 0.11 mg/kg, 0.22 ± 0.04 mg/kg and 0.174 ± 0.05 mg/kg respectively. Of the two toxic metals, lead concentrations are very high in all three organs of the complex during low water.

**Figure 4:** Concentration of lead and cadmium in fish organs

As for the sum of lead and cadmium concentrations in the various fish organs (Figure 5), it should be noted that lead was significantly higher during low water than during high water ($p=0.014$), and fish are more polluted with lead than with cadmium. Results at water level show that the concentration was higher in water during high water, whereas these pollutants are more concentrated in fish during low water.




Figure 5: Sum of lead and cadmium concentrations in whole fish (mg/Kg)

3.4 The bioaccumulation factor for lead and cadmium in fish

The bioaccumulation factor for lead and cadmium in *Sarotherodon melanotheron* fish is presented in Table 3. The concentration of lead in fish was 14.71 to 382.56 times higher than in water, and the concentration of cadmium in fish was 26.93 to 34.52 times higher than in water.

Table 3: Bioaccumulation factor for lead and cadmium in whole fish

Sites	Seasons	Heavy metals	Water concentration	Bioaccumulation factor
Lake Nokoué	High water	Cadmium	$0,017 \pm 0,01$	$26,93 \pm 7,23$
		Lead	$0,054 \pm 0,07$	$14,71 \pm 6,72$
	Low water	Cadmium	$0,010 \pm 0,001$	$31,35 \pm 0,71$
		Lead	$0,070 \pm 0,04$	$382,56 \pm 240,12$
Porto-Novo Lagoon	High water	Cadmium	$0,011 \pm 0,003$	$29,75 \pm 13,33$
		Lead	$0,141 \pm 0,11$	$28,78 \pm 13,08$
	Low water	Cadmium	$0,010 \pm 0,0005$	$34,52 \pm 11$
		Lead	$0,070 \pm 0,04$	$262,79 \pm 65,5$

3.4. Correlation matrix between lead, cadmium in whole fish and physico-chemical parameters

The correlation coefficients between physico-chemical parameters and toxic metals in whole fish (Table 4) shows that cadmium is strongly negatively correlated with dissolved oxygen and pH. Oxygen and pH are also positively correlated. But there is no correlation between lead in fish and the physico-chemical parameters measured.

Table 4: Correlation matrix between lead and cadmium in fish and physico-chemical parameters

Variables	r et p	Salinity	Dissolved ox- ygen (mg/l)	T°C	pH	Lead in fish	Cadmium in fish
Salinity	r	1					
	p	0					
Dissolved oxygen (mg/l)	r	-0,13	1				
	p	0,79	0				
T °C	r	0,68	-0,285	1			
	p	0,13	0,584	0			
pH	r	0,07	0,864	-0,385	1		
	p	0,88	0,026	0,451	0		
Lead in fish	r	0,59	0,170	0,506	0,089	1	
	p	0,21	0,747	0,305	0,867	0	
Cadmium fish	r	0,46	-0,820	0,620	-0,816	0,361	1
	p	0,35	0,046	0,189	0,047	0,482	0

“r” represents the coefficients of determination and ‘p’ represents p-values

3.5 Health risks associated with eating this fish

The Weekly Exposure Doses (WED) was assessed 15 to 90 kg using the reference of the World Health Organisation (WHO), which is 0.025 for lead and 0.007 for cadmium. The WED of the lowest weight up to 90 kg exceeded the Weekly Maximum Tolerable Dose (WMTD) set by the WHO for lead. This reveals that anyone weighing between 0 and 90 kg who consumes around 400 g of this fish per week is exposed to a major health risk linked to lead. As for cadmium, the WED up to 9 kg was exactly at the limit of the MLD given by the WHO (figure 7). This shows that the health risk for eating this fish associated with cadmium is lower




Table 5: Weekly Exposure Dose to lead and cadmium

4. DISCUSSION

Physico-chemical parameters

The physico-chemical parameters of an aquatic environment are basic factors for the survival and growth of aquatic organisms (Zandagba *et al.*, 2016). The lowest mean temperature obtained was $29.02 \pm 6.27^{\circ}\text{C}$ during high water, and the highest was $33.41 \pm 0.74^{\circ}\text{C}$ during low water. These values exceed those obtained by Capo-chichi *et al.* (2022), which were 26.80°C and 32.40°C , as well as (Agbohessi *et al.*, 2023b) and (Lalèyè *et al.*, 2022) in Lake Nokoué-lagune at Porto-Novo. These values also exceed the standard set by Water Quality Evaluation System (SEQ-Eau) (SEQ-Eau, 2003). This observed difference could be explained by several reasons, among which we can list the increase in the degree of organic pollution in this ecosystem and at our sampling sites (Capo-chichi *et al.*, 2022). Global warming could also be a cause of rising temperatures in aquatic environments. Hani *et al.* (2019) have shown that in an aquatic environment, the decrease in fish reproductive capacity was also linked to the increase in temperature (Agius and Robert, 2003).

Dissolved oxygen levels range from 2.11 to 5.38 mg/l, below the value set by SEQ-Eau (2003) for very good water quality. This value (2.11 mg/l) is below the values obtained by Capo-chichi *et al.* (2022) in Lake Nokoué. Dissolved oxygen in aquatic environments ensures fish respiration. A decrease in the oxygen content of an aquatic environment can be due to a rise in temperature, decomposition of organic matter and low water flow (Stantec, 2015). The high temperature recorded for this study may support this low oxygen level. The complex also faces heavy pollution of all kinds, even organic (Agbandou *et al.*, 2018), with the Acadja practice occupying 60% of the surface area of Lake Nokoué, which requires excessive use of wood that is then abandoned in the water. This helps reduce the flow of water in the complex, which also reduces oxygen levels. A drop in oxygen levels could cause mortality or reduce fish growth. The pH ranges from 6.21 to 8.62. These

values are close to those found by several other authors on Lake Nokoué (Capo-chichi *et al.*, 2022; Agbohessi *et al.*, 2023b). These values are also within the standards set by SEQ-Eau (2003).

Heavy metals in water and sediment

The average concentration of lead recorded in the water over the two seasons exceeded the standards set by the SEQ-Eau, which is 30 µg/l (SEQ-Eau, 2003). These values are close to those found by several authors on the Lake Nokoué - Porto-Novo lagoon complex (Kaki *et al.*, 2011; Agbandou *et al.*, 2018; Elegbede Manou *et al.*, 2020; Agbohessi *et al.*, 2023c). The same observations were made for the concentration in the sediment and that of cadmium found in the water and sediment. Exposure of fish to cadmium induces histopathological alterations in the liver, gills and kidney (Pantung *et al.*, 2008) and many other molecular and biological alterations. This high concentration is due to the activities carried out around and on the complex. Benin's largest brewing industry, which produces and stores alcoholic beverages, is located near Lake Nokoué. The high concentration of lead can also be explained by the proximity of these industries, which are also sources of lead (Kayalto and Mbofung, 2013). The results of these studies revealed that the concentration of both heavy metals is highest in water and sediments during floods. The same observations were made by Babalola and Fiogbe (2016) in the same ecosystem, particularly in the Porto-Novo lagoon. The high water levels correspond to the most intense rainy seasons of the year, and the complex is fed by several tributaries from northern Benin (Colleuil and Texier, 1987). In addition, Lake Nokoué receives all the rainwater runoff from the surrounding towns, the town's industries, numerous household waste disposal sites, and the many markets in the area due to its geographical location. Around 80% of Benin's industries are located along this waterway (Agbandou *et al.*, 2018). Industrial discharges are a potential source of heavy metals, particularly lead and cadmium (Angerville, 2009; Kayalto and Mbofung, 2013; Cuniasse and Glass, 2020). During rainfall, this complex also receives all urban discharges from the rainwater collectors of Cotonou and Abomey-Calavi (the most populated and industrialized cities in Benin) (Mama, 2010) which are a major source of heavy metals (lead and cadmium) at concentrations 7 to 26 times higher than in their areas of origin (Verge and Petit, 2009). The Lake Nokoué-Porto-Novo lagoon complex is also Benin's main transport route for petroleum products, particularly gasoline, and receives these products on a daily basis (Dovonou, 2019). These products are sources of lead (Loustau Cazalet, 2012). The high rain season is a period when the transport of petroleum products on the complex is more difficult, as high water and strong currents cause the boats transporting these products to capsize. These petroleum products are more likely to be spilled in the complex during the rainy season because they are sold not only on the shore but also in the lagoon complex, particularly in lakeside villages such as Ganvié, Sô-ava, Dékanmey, and Houédogbadji. They are sometimes accidentally spilled into the ecosystem due to leaks or explosions of cans during transport, or deliberately spilled by transporters to ensure the balance of boats or to escape customs or security checks (Agbohessi, Guedegba, *et al.*, 2023). All this explains the high concentrations of lead and cadmium recorded during high water in this ecosystem.

Heavy metal contamination of ichthyological fauna

The heavy metals present in this aquatic environment are more concentrated in fish than in water and sediment. Fish receive a large proportion of the heavy metals drained into this ecosystem. The calculated bioaccumulation factor shows that the amount accumulated by fish is as high as 382.56 in water mainly in low water. Studies conducted by Babalola (2015) in the same ecosystem in 2014 and 2015 showed that the species *Sarotherodon melanotheron* exhibited bioaccumulation of lead and cadmium 214.92 times and 23.18 times higher than their concentration in water, respectively. According to several studies conducted on the ecosystem between 2001 and 2023, an increase in lead and cadmium concentrations has been observed, and these heavy metals are the most studied in the environment (Chouti *et al.*, 2010; Kaki *et al.*, 2011; Hounkpatin *et al.*, 2012; Yehouenou *et al.*, 2013; Babalola and Fiogbe, 2016; Chouti *et al.*, 2017, 2020; Vodougnon *et al.*, 2018; Azon *et al.*, 2021; Agbohessi *et al.*, 2023b). This shows that the ecosystem is chronically exposed to these heavy metals. This explains the high bioaccumulation of lead in this species in 2024. Studies conducted by Agbohessi *et al.* (2023b) have also shown that the liver tissue of this species sampled in the same ecosystem exhibits severe histopathological alterations, particularly cellular necrosis. This could explain why their livers can't get rid of the heavy metals that have built up, which is why we've seen these pollutants accumulate. In the Lagos Lagoon in Nigeria, despite the fact that the concentration of heavy metals assessed was low and below standards in the water, *Sarotherodon melanotheron* fish caught in this lagoon accumulated these heavy metals in concentrations exceeding those found in the water and the recommended standards (Omoregham Jumoke and Okeke Ebele, 2025). Studies have also shown that the bioaccumulation factor for lead assessed in certain fish species (*Tilapia zillii*, *Chrysichthys nigrodigitatus*, *Synodontis membranaceus*) in the lower Volta basin in Ghana exceeded several hundred to over 1,000, indicating active and worrying bioaccumulation (Biney, 1991). Similarly, in the Ogun River in Nigeria, studies on *Clarias gariepinus* revealed very high concentrations of lead in the muscles (17.45 mg/kg) (Adeboyejo *et al.*, 2015). This shows that in ecosystems polluted with heavy metals, lead also appears to have a high capacity for bioaccumulation in fish muscle tissue. Lake Nokoué, a lagoon in Porto-Novo, appears to be one of the most lead-polluted aquatic ecosystems in West Africa.

Toxic metals are endocrine disruptors, they can cause feminization, decrease the production of 17 β -estradiol (E2), inhibit the synthesis of estradiol, and inhibit the effects of androgens (Scholz and Mayer, 2008). According to ecotoxicological studies, cadmium negatively affects the hypothalamic-pituitary-gonadal-hepatic axis, estradiol, and vitellogenin, even at very low concentrations of cadmium in water (5-10 μ g/l) (Hachfi, 2013). Exposure of fish to low concentrations of lead nitrate (0.1, 1, 3, and 10 μ g/L) for 24 hours resulted in inhibition of steroid production (Singh *et al.*, 2008). While the concentrations observed in the water of this ecosystem exceed those reported in these studies. These metals also cause histopathological alterations and malformations in embryos (Osman *et al.*, 2007). They also affect the biochemical functions of the kidneys and liver of fish (Elarabany and Bahnasawy, 2019) (Elarabany and Bahnasawy, 2019).

The muscles, gills, and gonads are the parts of the fish that are consumed. These organs have accumulated up to 16.41 mg/kg of metals, whereas the limit set by the FAO/WHO in “CODEX STAN 193-1995” and Seventy-third meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA) (JECFA, 2011) is 0.2 mg/kg for lead and 0.01 mg/kg for cadmium. The study revealed that during the dry season, when water levels are low, the concentration of heavy metals in fish is significantly higher, whereas the concentration in water is high during high water. This clearly explains the phenomenon of bioaccumulation. After pollutants are released into the ecosystem during high water, it takes some time for fish to accumulate them, and this accumulation becomes apparent during periods of low water. This absorption occurs directly through the respiratory system via the gills (Omoregham Jumoke and Okeke Ebele, 2025), which shows a high concentration in the gills. Absorption also occurs through the food chain when aquatic plants and micro- and macro-invertebrates that come into contact with the sediment are consumed (Soltani *et al.*, 2019). This concentration found in fish exceeds that obtained by several authors on the same fish species in the same ecosystem (Youssao *et al.*, 2011; Goussanou, Youssao, *et al.*, 2018; Elegbede Manou *et al.*, 2020; Agbohessi, Onibon, *et al.*, 2023b). This complex is more polluted with lead than the Anambra River in Nigeria, where *Clarias gariepinus* and *Oreochromis niloticus* fish are believed to be contaminated with toxic metals, with accumulated lead concentrations of 1,062 ppm (Ogbuene *et al.*, 2024). Appropriate measures must be taken to limit sources of pollution and consider a plan to clean up this ecosystem.

The Weekly Exposure Doses (WED) assessed for lead show that anyone weighing between 0 and 90 kg and consuming around 400 g of this fish per week is at high risk of Lead poisoning. As for cadmium, WED up to 9 kg are exactly at the limit of the Maximum Tolerable Weekly Dose (MTHWD) given by the WHO. It was shown that consumption of 150 g of fish from Lake Nokoué per day presented a lead poisoning risk for a 30 kg child (Elegbede Manou *et al.*, 2020). Whereas in the present study, the lead related poisoning dose assessed is less than half that found by this author for a weight range of 0 to 90 Kg. This shows that the level of pollution in this complex is very high, and the population is exposed to great danger. The population needs to be aware of the risk of poisoning linked to the consumption of fish from the complex. A toxicology study is needed to assess the health status of all fish products from this ecosystem destined for consumption. Chronic exposure of the human body to lead has numerous health drawbacks, such as reproductive disruption, cancer, hypertension, renal failure, neuropathy and hematological effects, etc. (Bismuth *et al.*, 2002). A study on the treatment of urban discharges from stormwater collectors in the communes that this ecosystem serves should be considered. It is also necessary to consider depolluting this watercourse to save the population that depends on it.

The influence of physico-chemical parameters on the bioaccumulation of heavy metals

cadmium concentrations in fish are negatively correlated with pH and dissolved oxygen. This negative correlation shows that cadmium is more mobile to accumulate in fish when pH and dissolved oxygen are low. According to the Canadian Environmental Protection Act, cadmium is more toxic and mobile in low pH eco-

systems (LCPE, 2009). The lake Nokoué-Porto-Novo lagoon complex is exposed to organic and chemical pollution (Agbohessi, Guedegba, *et al.*, 2023c), affecting physico-chemical parameters (Capo-chichi *et al.*, 2022). In the present work, the recorded oxygen level was low and the pH was also low at several sites. This could facilitate cadmium accumulation in fish and cause more damage to the species' physiology. All these phenomena expose the ecosystem to imminent degradation if no solution is envisaged.

A study of the physiological state and, above all, the reproductive system of the fish is needed to determine the damage this pollution would have caused to the physiology and sustainability of the aquatic fauna.

5. CONCLUSIONS

Our study confirmed that the complex is polluted with heavy metals, in particular lead and cadmium, through their accumulation in water, sediment and aquatic fauna. This poses a risk of metal poisoning when these fish are consumed by the general public. Appropriate measures to limit sources of pollutants must be taken. Consider an alternative route for transporting petroleum products, or opt for a safer means of transport, which are also a major source of lead.

Author Contributions

- Gisèle HOUEDJISSI: Conceptualization and conservation of data and Writing - original draft, Writing;
- Maxime Machioud Sangare-Oumar: Validation, Writing - revision;
- Prudencio Agbohessi: Writing - editing;
- Chabi Oloushegun Onibon: Sample collection;
- Jean-Michel AKPO: Supervision.

Funding

This research was funded by the African Center of Excellence for Water and Sanitation (C2EA) of the National Water Institute (INE) of the University of Abomey Calavi.

Acknowledgments

The authors sincerely thank Dr Alexis HOUNDJI, Dr Abdoulaye ISSOTINA ZIBRILA, Dr Besmer Régis AHISSOU, Mr Eric GLODJO and Mr Condé-Abalou Roland BAMAI for their sacrifices in carrying out this work

Conflicts of Interest:

The authors declare no conflicts of interest.

REFERENCES

1. Adeboyejo, A.O., Morenikeji, O.A. and Fagade, S.O., 2015. Assessment of Heavy Metals in Fish Species from Ogun River, Nigeria. *Journal of Environmental and Occupational Science*, 4(1)

2. Agbandou, B., Houessou, D., Thoto, F. and Chabi, G., 2018. *Déséquilibre écosystémique du complexe lagunaire lac Nokoué – lagune de Porto-Novo. Note politique ACED*. Abomey-Calavi. <https://www.aced-benin.org/fr/publications>.
3. Agbohessi, P., Guedegba, L., Pelebe, R. and Toko, I.I., 2023a. Exposition de la faune ichthyenne et autres animaux aquatiques aux polluants organiques , chimiques et biologiques dans le lac Nokoué au Sud du Bénin : Synthèse documentaire. *Revue Marocaine des Sciences Agronomiques et Vétérinaires*, 11, pp. 360–372.
4. Agbohessi, P., Kinsicounon, G., Ouedraogo, A., Guedegba, L., Houndji, A. and Imorou Toko, I., 2023c. Changes in blood tissue of estuarine tilapia *Sarotherodon melanotheron* (Rüppell , 1852) captured in polluted Lake Nokoué (Benin , West Africa). *Research Journal of Animal, Veterinary and Fishery Sciences*, 11(1), pp. 1–7.
5. Agbohessi, P., Onibon, C., Pelebe, E.O.R., Mattah, D.A.P. and Guedegba, Nicresse Léa Imorou Toko, Ibrahim Chikou, A., 2023b. Assessment of the health status of Lake Nokoué (Benin, West Africa) exposed to multiple pollutants during flood season. *Chemistry and Ecology*, pp. 1–27. <https://doi.org/10.1080/02757540.2023.2253225>.
6. Agius, C. and Robert, R.J., 2003. Melano-macrophage centres and their role in fish pathology. *Journal of Fish Diseases*, 26, pp. 499–509. <https://doi.org/10.2174/157018081503180131102747>.
7. Angerville, R., 2009. *Evaluation des risques écotoxicologiques liés au déversement de Rejets Urbains par Temps de Pluie (RUTP) dans les cours d ' eau : Application à une ville française et à une ville haïtienne. Thèse de doctorat*.
8. Azon, N.B.N., Hounkpe, P., Adoukpe, J.G., Degila, H.W. and Aina, M.P., 2021. Direct Mercury Analyzer (DMA) determination of mercury distribution in the sediments of lake Nokoué in Benin Republic-West Africa. *International Journal of Biological and Chemical Sciences*, 15(1), pp. 306–316. Available at: <https://doi.org/10.4314/ijbcs.v15i1.27>.
9. Babalola, O.A., 2015. *Evaluation of metal pollutants in water, sediments and four economically important fish specie from Porto-Novo lagoon, Republic of Benin. Thèse de Doctorat*. Université d'Abomey-Calavi.
10. Babalola, O.A. and Fiogbe, D.E., 2016. Seasonal Variation Assessment and Correlation Coefficient of Metal Pollutants in Sediments and Water from Porto Novo Lagoon Ecosystem, Benin Republic. *American Journal of Educational Research*, 4(12), pp. 976–982. Available at: <https://doi.org/10.12691/education-4-13-10>.
11. Barbier, M.G., 2011. *Rapport sur les perturbateurs endocriniens, le temps de la précaution*.
12. Başıyigit, B. and Tekin-özan, S., 2013. Concentrations of Some Heavy Metals in Water , Sediment , and Tissues of Pikeperch (*Sander lucioperca*) from Karataş Lake Related to Physico-Chemical Parameters , Fish Size , and Seasons. *Pol. J. Environ. Stud.*, 22(3), pp. 633–644.
13. Biney, C.A. (Conseil de recherches scientifiques et industrielles du G.-C., 1991. Bioaccumulation of Heavy Metals in Fish Species of the Lower Volta Basin, Ghana. *Environmental Health Perspectives*, 103(1).
14. Bismuth, C., Baud, F., Conso, F., Dally, S., Fréjaville, J.-P., Garnier, R. and Aeger, J.A., 2002. *Les effets du plomb sur la santé. Rapport DGS/Sous-direction de la gestion des risques des milieux Bureau*. France.
15. Capo-chichi, H.B.P., Adandedjan, D., Agblonon Houelome, M. and Lalèyè, P., 2022. Physico-chimie et pollution organique du lac Nokoué au Sud du Bénin . *Journal of Applied Biosciences* 170:, 170, pp. 17752–17774. <https://doi.org/10.35759/JABs.170.7>.
16. Chouti, W., Agonzan, S., Tometin, A.L. and Mama, D., 2017. Étude de la spéciation chimique du plomb dans les sédiments de la lagune de Porto-Novo, Sud Bénin. *Afrique Science*, 13(3), pp. 193–205.

17. Chouti, W., Hounkanrin, F., Adandedji, F.M. and Mama, D., 2020. Chemical characterization of metallic trace elements in aquatic plants : A case study of some plants in Porto Novo lagoon , South Benin. *International Research Journal of Environmental Sciences*, 9(4), pp. 66–73.
18. Chouti, W., Mama, D. and Alapini, F., 2010. Etude des variations spatio-temporelles de la pollution des eaux de la lagune de Porto-Novo (sud Bénin). *International Journal of Biological and chemical Sciences*, 4(4), pp. 1017–1029.
19. Colleuil, B. and Texier, H., 1987. Le complexe lagunaire du Lac Nokoué et de la lagune de Porto-Novo (Bénin). *Institut Français de Recherche Scientifique pour le Développement en Coopération* [Preprint], (13).
20. Cuniasse, B. and Glass, T., 2020. *Métaux lourds. Rapport Secten - Citepa. Émissions de gaz à effet de serre et de polluants atmosphériques en France*. France. citepa.org/fr/secten.
21. Dovonou, F., 2019. Impacts des Hydrocarbures Aromatiques Polycycliques sur les poissons: Cas des Tilapias du lac Nokoué au Sud du Bénin (Afrique de l'Ouest). *European Scientific Journal*, 15(36), pp. 458–474. <https://doi.org/10.19044/esj.2019.v15n36p458>.
22. Elarabany, N. and Bahnasawy, M., 2019. Comparative and Interactive Biochemical Effects of Sub-Lethal Concentrations of Cadmium and Lead on Some Tissues of the African Catfish (*Clarias gariepinus*). *Toxicological Research*, 35(3), pp. 249–255. Available at: <https://doi.org/10.5487/TR.2019.35.3.249>
23. Elegbede Manou, B., Sohounnon, M., Dovonou, F., Koumolou, L. and Etorh, P.A., 2020. Évaluation des risques d'intoxication alimentaire liés à la consommation des produits halieutiques du lac Nokoué , Bénin , Afrique de l ' Ouest Résumé. *Afrique SCIENCE*, 16(5), pp. 292–302.
24. Goussanou, A., Youssao, A., Karim, A., Toleba, S.S., Dagan, B.S., Gnandi, K. and Youssao, I., 2018. *Cardisoma armatum* (Herklots, 1851) dans le complexe lac Nokoué-lagune de Porto-Novo au Sud Bénin. *Afrique SCIENCE*, 14(2), pp. 255–266. <http://www.afriquescience.info>.
25. Hachfi, L., 2013. *Effets de stress physico-chimiques sur l'expression de gènes impliqués dans l'homéostasie et la fonction de l'axe hypothalamo-hypophyso-gonado-hépatique chez le loup (Dicentrarchus labrax L.)*. Thèse présentée pour obtenir le grade de docteur en Science de l. Université de Toulon.
26. Hani, I.M.Y., Turies, C., Palluel, O., Delahaut, L., Bado-Nilles, A., Geffard, A., Dedourge-Geffard, O. and Porcher, J.-M., 2019. Effects of a chronic exposure to different water temperatures and / or to an environmental cadmium concentration on the reproduction of the threespine stickleback (*Gasterosteus aculeatus*). *Ecotoxicology and Environmental Safety*, 174, pp. 48–57. <https://doi.org/10.1016/j.ecoenv.2019.02.032>.
27. Hounkpatin, A.S., Etorh, A.S., Sèzonlin, M., Guédénon, P., Elégbédé, B., Boni, G., Dougnon, V., Montcho, S., Kéké, E. and Boko, M., 2012. Pollution of aquatic ecosystems by heavy metals at Ganvié's lacustrine city (Benin). *International Research Journal of Biotechnology*, 3(6), pp. 81–87.
28. INSAE (2015) *RGPH4 : QUE RETENIR DES EFFECTIFS DE POPULATION EN 2013 ?*
29. JECFA (Seventy-third meeting of the Joint FAO/WHO Expert Committee on Food Additives), 2011. Series No. 64. Geneva, Switzerland, p. 543.
30. Kaki, C., Patient, G., Nelly, K., Etorh, P.A. and Adechina, R., 2011. Evaluation of heavy metals pollution of Nokoue Lake. *African Journal of Environmental Science and Technology*, 5(3), pp. 255–261.
31. Kayalto, B. and Mbofung, C.M.F., 2013. Contribution à l'évaluation de la contamination par les métaux lourds, de trois espèces de poissons, des sédiments et des eaux du lac Tchad. *HAL Open Science*, p. 98.

32. Lalèyè, K.R., Agadjihouèdé, H., Lédèroun, D., Houélomè, T.A., Chikou, A. and Lalèyè, P., 2022. Impact de l'exploitation du sable fluvio-lagunaire sur la qualité physico-chimique de l'eau des écosystèmes aquatiques du sud-Bénin. *Biotechnologie, Agronomie, Société et Environnement*, 26(2), pp. 121–135.
33. Lalèyè, P., Chikou, A., Philippart, J., Teugels, G. and Vandewalle, P., 2004. Étude de la diversité ichthyologique du bassin du fleuve Ouémé au Bénin (Afrique de l' Ouest). *Cybium*, 28(1), pp. 329–339.
34. Lalèyè, P., Philippart, J.-C. and Poncin, P., 1995. Biologie de la reproduction de deux espèces de Cbrysicbtbys (Siluriforme, Bagridae) du lac Nokoué et de la lagune de Porto-Novo au Bénin. *African Zoology*, 109, pp. 213–224.
35. LCPE, 2009. *Loi Canadienne sur la Protection de l'Environnement (LCPE). Le cadmium et ses composés. Liste des substances d'intérêt prioritaire. Rapport d'évaluation.*
36. Loustau Cazalet, M., 2012. *Caractérisation physico-chimique d'un sédiment marin traité aux liants hydrauliques – Évaluation de la mobilité potentielle des polluants inorganiques. Thèse présentée pour l'obtenir le grade de docteur.*
37. Mama, D., 2010. *Méthodologie et résultats du diagnostic de l'eutrophisation du lac Nokoué (Bénin). Thèse de doctorat.* Université de Limoges.
38. Omoregham Jumoke, K. and Okeke Ebele, G., 2025. Heavy metal concentration in Sarotherodon melanotheron (blackchin tilapia) from the Lagos lagoon complex, and the potential human health risk from consumption. *LASU Postgraduate School Journal*, (July 2024), pp. 162–176. Available at: <https://doi.org/10.5281/zenodo.14481937>.
39. Osman, A.G.M., Wuertz, S., Mekki, I.A., Exner, H.-J. and Kirschbaum, F., 2007. Malformations induites par le plomb dans les embryons du poisson-chat africain *Clarias gariepinus* (Burchell, 1822). *Wiley InterScience*, 22, pp. 375–389. Available at: <https://doi.org/10.1002/tox.20272>.
40. OMS, 2017. *Organisation Mondiale de la Santé. Directives de qualité pour l'eau de boisson. Quatrième édition.*
41. Pantung, N., Helander, K.G., Helander, H.F. and Voravit, C., 2008. Histopathological alterations in hybrid walking catfish (*Clarias macrocephalus* x *Clarias gariepinus*) during acute and subacute exposure to Nuntiya cadmium. *EnvironnementAS*, 1, pp. 22–27.
42. Rossi, N., 2009. Écologie des communautés planctoniques méditerranéennes et étude des métaux lourds (Cuivre, Plomb, Cadmium) dans différents compartiments de deux écosystèmes côtiers (Toulon, France). Thèse de doctorat pour l'obtention du grade de docteur. *HAL Open Science*, p. 287.
43. Rodier, J., Legube, B., Merlet, N. and Al., E., 2009. Assessment of Ground Water Quality in Baba I Village, North-West Cameroon. *Journal of Geoscience and Environment Protection* [Preprint].
44. Scholz, S. and Mayer, I., 2008. Molecular biomarkers of endocrine disruption in small model fish. *HAL Open Science*, 293(57). Available at: <https://doi.org/10.1016/j.mce.2008.06.008>.
45. SEQ-Eau, 2003. *Système d'évaluation de la qualité de l'eau des cours d'eau. Rapport de présentation version 2.*
46. Singh, J., Kant, K., Sharma, H.B. and Rana, K.S., 2008. Bioaccumulation of Cadmium in Tissues of *Cirrhina mrigala* and *Catla catla*. *Asian Journal of Experimental Sciences*, 22(3), pp. 411–414.
47. Soltani, N., Moore, F., Keshavarzi, B., Sorooshian, A. and Javid, R., 2019. Potentially toxic elements (PTEs) and polycyclic aromatic hydrocarbons (PAHs) in fish and prawn in the Persian Gulf, Iran. *Ecotoxicology and Environmental Safety*, pp. 251–265. Available at: <https://doi.org/10.1016/j.ecoenv.2019.02.005>.
48. Stantec, 2015. *Projet Sisson: Rapport final d'évaluation de l'impact sur l'environnement.*
49. Verge, G. and Petit, C., 2009. Contamination des organismes marins du littoral français par le Cadmium : évaluation et gestion du risque. *Revue Médecine Vétérinaire*, 160(10), pp. 484–490.

-
50. Vodougnon, H., Lederoun, D., Amoussou, G., Adjibogoun, D. and Laleye, P.A., 2018. Ecologic stress in fish population of Lake Nokoué and Porto-Novo Lagoon in Benin. *International Journal of Fisheries and Aquatic Studies*, 6(3), pp. 292–300. Available at: <https://www.fisheriesjournal.com/archives/2018/vol6issue3/PartD/6-2-41-728.pdf>.
 51. Yehouenou, E.A.P., Adamou, R., Azehoun, P.J., Edoth, P.A. and Ahoyo, P., 2013. Monitoring of Heavy Metals in the complex “Nokoué lake - Cotonou and Porto-Novo lagoon” ecosystem during three years in the Republic of Benin. *Research Journal of Chemical Sciences*, 3(5), pp. 12–18. Available at: www.isca.in.
 52. Youssao, A., Soclo, H.H., Bonou, C., Vianou, K., Gbaguidi, M. and Dovonou, L., 2011. Evaluation de la contamination de la faune ichthyenne dans le complexe lagunaire Nokoué – chenai de Cotonou par le plomb : cas des espèces *Sarotherodon melanothron*, *Tilapia guineensis* et *Hemichromis fasciatus* (Bénin). *International Journal of Biological and chemical Sciences*, 5(April), pp. 595–602.
 53. Zandagba, J., Adandedji, F.M., Mama, D., Chabi, A. and Afouda, A., 2016. Assessment of the Physico-Chemical Pollution of a Water Body in a Perspective of Integrated Water Resource Management: Case Study of Nokoué Lake. *Journal of Environmental Protection*, 7, pp. 656–669. <https://doi.org/10.4236/jep.2016.75059>.