Prepublished copy Pollution Technology

Original Research Paper

Health Risk Assessment Among Biogas and Conventional Cooking Fuel Users in Different Socioeconomic Conditions of Rural West Bengal

Tapas Bagdi¹, Apurba Koley², Niladri Das³, Suman Das², Sujan Gayen², Anuradha Sarkar¹, Sreya Ghosh¹, Sushil Kumar⁴, Shibani Chaudhury², Amit Kumar Hazra¹, Andrew B Ross⁵, Deep Chakraborty⁶†, Srinivasan Balachandran²

- ¹ Social Energy Laboratory, Department of Lifelong Learning and Extension, Rural Extension Centre, Institute of Rural Reconstruction, Visva-Bharati, Sriniketan, West-Bengal, India 731236
- ² Bio-Energy Laboratory, Department of Environmental Studies, Siksha-Bhavana, Visva-Bharati 731235
- ³ Department of Geography, Hiralal Bhakat College, Nalhati, Birbhum, West-Bengal
- ⁴ Environmental Chemistry Laboratory, Department of Environmental Studies, Siksha-Bhavana, Visva-Bharati 731235
- ⁵ School of Chemical and Process Engineering, University of Leeds, Leeds, LS2 9JT, United Kingdom
- ⁶ Department of Environmental Health Engineering, Sri Ramachandra Faculty of Public Health, Sri Ramachandra Institute of Higher Education and Research (Deemed to be University), Porur, Chennai-600116

[†]Corresponding author: Deep Chakraborty (<u>deepckbty@gmail.com</u>)

<u>Orcid: 0000-0002-0513-8100</u> Orcid: 0000-0003-4247-408X)

Key Words	Household air pollution, Exposure load pattern, Health risk assessment,
	biogas, Monte-Carlo simulation model
DOI	https://doi.org/10.46488/NEPT.2025.v24i04.B4295 (DOI will be active only after
	the final publication of the paper)
Citation of the	Bagdi, T., Koley, A., Das, N., Das, S., Gayen, S., Sarkar, A., Ghosh, S., Kumar,
Paper	S., Chaudhury, S., Hazra, A. K., Ross, A. B., Chakraborty, D., Balachandran,
	S., 2025. Health risk assessment among biogas and conventional cooking fuel
	users in different socioeconomic conditions of rural West Bengal. Nature
	Environment and Pollution Technology, 24(4), p.B4295.
	https://doi.org/10.46488/NEPT.2025.v24i04.B4295

Abstract: In many rural areas of developing countries, solid fuel use is still widespread. The present study aims to investigate the Household Air Pollution (HAP) exposure effect from traditional biomass fuels and biogas on the health of rural women. The results revealed that nearly 93% of rural families utilize conventional fuels for daily cooking and heating purposes, whereas clean fuels like biogas users were minimal. However, high-income, educated, elderly, well-structured houses (*Pucca*), and hierarchically more advanced families were observed to cook with biogas fuel. Further, the present study also used spatial HAP mapping and land use mapping models to analyze exposure load patterns and sources of solid fuel availability, respectively. The clean fuel biogas burning showed the lowest HAP concentration compared to traditional fuels except for NO₂ (1.14 ± 0.05 ppm), which also represents in case of health risk estimation. The biogas users also observed the lowest COHb% (0.008 ± 0.01) than conventional fuels. The health risks associated with SO₂ and NO₂ for biogas users were revealed to be lowest in both acute and chronic instances. Monte-Carlo probabilistic model observed that coal cake may pose high health risks among traditional fuels when concerning PM_{2.5}, SO₂, and NO₂, but in the case of COHb %, cow dung users showed the highest health risk (0.39 ± 0.02). These findings have significant implications for public health, suggesting that promoting the use of cleaner cooking fuels, particularly biogas, found less affected by muscular pain and eye irritation, than biomass users, could lead to substantial health benefits

for rural populations. This finding also indicates that government intervention should be required to enhance the utilization of cleaner cooking fuels in rural India for women's safety.

1. INTRODUCTION

Household air pollution (HAP) is the eighth most significant contributor to the global disease burden, and growing concern is caused by the combustion of solid fuels that pose a significant threat to human health (Manuel and Gautam, 2023). About 2.6 billion people worldwide still cook with solid, unprocessed fuels (Mitra et al. 2023), making them vulnerable to the toxic effects of HAP. Recently, the World Health Organization (WHO) reported the HAP was responsible for a distressing global annual death of 3.2 million, including approximately 237,000 child fatalities in 2020 (WHO 2022). This exposure to HAP becomes a cause of low birth weight, pregnancy complications (Amegah et al. 2024), acute respiratory infections (Enyew et al. 2021), respiratory impairment (Pathak et al. 2019), and cardiovascular disease (Mocumbi et al. 2019) and cognitive impairment (Krishnamoorthy et al. 2018), chronic obstructive pulmonary disease (COPD) (Wu et al. 2022). Most rural households mainly depend on solid biomass due to their lower socio-economic conditions, and women collect fuelwood for 5 to 8 hours a day, taking up 20% of their time and carrying 25–50 Kg of firewood for several miles (Matano 2022).

India, a country heavily dependent on agriculture, has a potential amount of biomass that may be processed using cutting-edge techniques to produce biogas energy. However, with the enormous emissions of greenhouse gases (GHGs) resulting from the combustion of fossil fuels, there is a growing need for alternative renewable fuels (Bagdi et al. 2023; Shane and Gheewala 2017). Methane gas yields about 802.29 KJ/mol of energy, while typical biogas (60% methane) produces 534.86 kJ/mol of energy (Mekonen et al. 2023). A small family-size biogas plant of one cubic meter capacity can save 0.43 kg of liquefied petroleum gas (LPG) daily (Aggarwal et al. 2021). In order to promote biogas technology in rural areas on large-scale applications for substitute cooking fuel, chemical fertilizer, and the generation of off-grid electricity, the government of India has initiated various initiatives, including the New National Biogas and Organic Manure Programme, Biogas Power Generation (offgrid) and Thermal Energy Application Programme, the Central Financial Assistant and the Sustainable Alternative Towards Affordable Transportation (Bharti 2019). The Ministry of New and Renewable Energy (MNRE) continues the National Bioenergy Programmed for FY 2021-22 to 2025-26, which recommended the implementation of NBP in two phases. Phase-I, approved with a budget of INR ₹ 8580 million, includes around USD 12.5 million for the Biogas Programme, supporting modest to medium-sized biogas plants. MNRE targeted the buildup 30795 number of small-scale biogas plants all over India by the financial year 2023-2024 (MNRE 2022). Despite the government of India's efforts to prioritize small-scale biogas plants in rural India, only 0.4% of rural Indian households use biogas as their primary energy source (Mottaleb 2019). According to a recent estimation in India, about 54% of households still rely on traditional solid biomass cooking fuels likewise firewood, cow dung cake, charcoal, and agricultural residues for cooking and heating purposes, which are prominent sources of indoor air pollution in developing nations (Sunil et al. 2021). Several scholarly investigations (Kumar et al. 2023; Shobande 2023; Zewdie et al. 2023) highlighted that the practice of burning solid fuels in inefficient stoves, attached with inadequate ventilation, can pose significant health risks. In addition, cooking with incomplete combustion of solid fuels, especially in unvented environments, can lead to elevated indoor levels of carbon monoxide (CO), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), particulate matter (PM_{2.5} and PM₁₀), volatile organic compounds (VOCs) and polyaromatic hydrocarbons (PAHs) (Churchill and Smyth 2021; Nawaz 2021).

Recently, India announced at COP26 the targets to achieve larger goals within the allotted period as per COP21. India set a 40% non-fossil fuel energy goal as part of the Paris Agreement to combat climate change (Sawhney, 2021). At the COP 26 summit in Glasgow, India's new objective has been revised to 500 gigawatts (GW) from renewable energy by the year 2030 to cut its carbon emissions by nearly a billion tones, aiming to

have a carbon-neutral economy by 2070 (Singh et al. 2023). The installed capacity of bio-power in India has reached 10.16 gigawatts, which will account for 10.12% of the country's total capacity for renewable energy sources by the end of 2021 (Kulyal and Jalal, 2022). Thus, there is a need to transform traditional energy into clean, modern, and sustainable energy to control climate change. Efficient, clean energy sources are linked to improved HAP, expanded opportunities for productive or recreational activities, and decreased dependence on biomass energy collection. In a research conducted by Afridi et al. (2023), it was observed that women spend an average of 23 hours per week on cooking activities.

Additionally, when cleaner fuel was used for cooking, there was an 18% reduction in cooking time. In their study, Waris and Antahal (2014) observed that the average cooking time per meal decreased from 1.08 to 0.75 hours when using biogas stoves. Anderman et al. (2015) discovered that women who utilize biogas spend an average of 40 minutes reduction cooking and 70 minutes reduction in collecting firewood per day compared to women in households that use solid biomass (Chowdhury et al. 2023). In order to mitigate the harmful effects of HAP and promote better health outcomes, various interventions have been proposed or tested. Among them, shifting towards cleaner cooking energy for example, biogas (Koley et al. 2022a), electricity, LPG (Bagdi et al. 2022) or solar power are used. Another approach involves using improved cookstoves that generate more energy while producing fewer smoke (Chakraborty et al. 2021). In addition, interventions may focus on enhancing ventilation and optimising kitchens' design, as well as promoting changes in cooking behaviours (Chakraborty et al. 2022).

The 68th round report by the National Sample Survey Organisation (NSSO) emphasised the significant dependence of rural families in India on traditional fuels. In 2011-12, it was found that 67.35% of rural house-holds relied on firewood and chips as their primary cooking fuel source. The substantial reliance on firewood highlights rural communities' challenges in getting cleaner and more efficient energy sources. Using firewood poses health hazards due to indoor air pollution and contributes to environmental degradation by causing deforestation. The results emphasize the pressing necessity to advocate for alternative, sustainable energy options such as biogas. Biogas has the potential to boost health outcomes, mitigate environmental consequences, and improve the standard of living in rural regions. Moreover, the availability of huge potential of biomass resources in the village like aquatic weeds (Koley et al. 2023; Koley et al. 2022b), algae (Koley et al. 2024b), agricultural residues, livestock resources (Dhungana et al. 2022; Bagdi et al. 2022), food waste (Nahar et al. 2024) etc. that may be used as feedstock for the biogas plant.

There are limited studies that have been conducted in the past on the comparison of health benefits from biogas over traditional cooking fuels under varying socio-economic conditions. This study investigated the relationship between socio-economic status, exposure load and land use patterns. Additionally, the study analyzed the health risk assessment among rural women who are using traditional and biogas fuels. This study highlights the health and socio-economic benefits of biogas over traditional fuels, assesses health risks among rural women, and explores exposure assessments and land use patterns to provide policy insights for promoting clean energy and sustainability.

2. Materials and Methods

2.1. Study Area

The present study was conducted in 'Sehalai' village (Lat 23.718238, Long 87.680435), located under Kasba gram panchayat in Bolpur-Sriniketan tehsils of Birbhum district, West Bengal, India (Fig. 1). The village covers 328.54 hectares of geographical area. As per the census (2011), the reported total population of this village is 1408 people with 315 households. In Sehalai village, out of the total population, 386 (27.41%) and 616 (43.75%) belong to SC and ST communities. The primary occupation of the villagers is agriculture, with

approximately 302.4 hectares of irrigated land and 2.4 hectares of forestland. Most villagers are engaged in agriculture (31%), while almost 60% rely on casual labour and other sectors. Sehalai village was chosen as the study region for this case study because it indicates rural Bengal, as reported by the National Family Health Survey (NFHS) 2018, which found that 71% of rural families in West Bengal use firewood as their major cooking fuel. This heavy dependency reflects the region's broader rural environment, providing common concerns such as health risks from indoor air pollution and degradation of the environment caused by deforestation. By focusing on Sehalai hamlet, the study intends to gain insight into these prevalent concerns and assess the possible benefits of implementing sustainable energy options such as biogas. This selection assures that Sehalai's findings and recommendations may be applied to similar rural settings in West Bengal, providing an extensive comprehension of energy usage dynamics and the need for cleaner, more efficient cooking solutions.



Fig. 1: Study Area (a) Birbhum District of West-Bengal; (b) Sehalai Village of Birbhum District (c) North Sehalai Village and (d) South Sehalai village

2.2. Personal Interview and Data Collection

The study region was selected on the following basic criteria: (i) The village is at least 3 km from the major road to reduce other pollution (noise, vehicular pollution); (ii) no air polluting industries are present in the surrounding areas; (iii) the study areas have knowledge about biogas; and (iv) respondents aged 18 to 69 years were chosen. The inclusion criteria were non-smoking women who cooked using solid fuels or biogas for at least 2 hours per day for the last 5 years, along with LPG as an emergency backup, and who had no chronic health conditions. Exclusion criteria include people who are entirely transitioning to LPG as their primary cooking fuel and ladies with chronic conditions.

The proposed research area's sample size is determined using a 95% confidential level and a 5% marginal error. According to the 2011 census report, the village has a total of 315 households. The proposed sample size is approximately 173 households out of the total number of households in the village. The recommended sample results are designed to ensure that they accurately represent the total population. The random sampling technique was employed to choose representative sample households from throughout the hamlet. The survey was conducted between July 2024 to September 2024. The questionnaire was originally developed in English and subsequently translated into Bengali, the native language of the study area. Participants were informed of the study's objective in Bengali, and informed consent was obtained prior to data collection. Participants who were literate provided written assent, while those who were unable to read or write confirmed their consent verbally. The authors surveyed to build rapport and enhance data gathering across the village. The questionnaire used a combination of open-ended, closed-ended, and multiple-response questions to collect both qualitative and quantitative data. A total of 173 households were conducted surveys through personal interviews to gather data on the households' demographic characteristics, such as the age of the head, head education, primary occupation of the head, average family income, daily cooking hours, cooking experience, kitchen and fuel type, agricultural land holding and housing structure. This information was collected using a standardised questionnaire with permission from the participating families and recorded necessary information.

2.3. Measurement of Household Air Pollution

A total of 173 households were surveyed to gain insights into their socio-economic conditions and fuel usage patterns. Among these 173 households, a subset of 20 were selected from two different colonies that primarily relied on wood, cow dung cake, coal cake, and biogas to measure HAP indoors. During the monitoring, 5 households were excluded from the study due to non-compliance with the research protocol. The HAP was measured using a real-time 8-channel Indoor Air Quality (IAQ) Monitor (YESAIR) developed by Critical Environment Technologies in British Columbia, Canada. The real-time monitoring has been carried out for 24 hours, including cooking and non-cooking periods in selected households. The devices were set up at a breathing-level height and 1.5 meters away from the stove (in a sitting position) (Chakraborty et al. 2014).

2.4. Quality Assurance/ Quality Control

Instruments were adequately charged and calibrated to maintain the data quality. The electrochemical sensors of CO (PNP-C), NO₂ (PNP-D), SO₂ (PNP-P), and CET-PM_{2.5} were calibrated before each sampling session using the supplied calibration tube attached with one end to the cylinder full of 100% Nitrogen. The instrument also has been standardized in a controlled environment with an observation of \pm 3% accuracy level.

2.5. Assessment of household air exposure and health risk

2.5.1. PM_{2.5} exposure

The degree of exposure was determined by Chakraborty and Mondal (2018), using the total concentration of $PM_{2.5}$ emitted the number of occupants, and the total amount of time spent in various circumstances while cooking. The exposure index was computed using Eq. 1. $PM_{2.5}$ guideline value taken from WHO (2021).

$$E_i = \sum_{i}^{l} \frac{C_i t_{ki}}{C_g t_a} \qquad \dots (1)$$

2.5.2. Intake concentration of PM_{2.5}

Mitra et al (2022) determined the intake concentration by calculating the average concentration of $PM_{2.5}$. Eq. 2 was utilized to calculate the identical value (USEPA, 1989).

$$IC = \frac{C_p \times E_t \times E_f \times E_d}{AT} \qquad \dots (2)$$

2.5.3. Intake concentration of PM_{2.5}

The majority of cardiovascular diseases are linked to $PM_{2.5}$ intake. As a result, the toxicological risk of $PM_{2.5}$ was assessed using the Hazard Quotient (HQ) using Eq. 3 (USEPA, 2005).

$$HQ = \frac{IC}{Rfc} \qquad \dots (3)$$

If the outcome values of HQ are more than 1, then there will be a chance to develop non-carcinogenic health risks among exposed groups.

2.5.4. Blood carboxyhemoglobin level assessment

Applying the approach proposed by Peterson and Stewart, the blood COHb% levels were estimated (Eq. 4) for each fuel burned in this study (Chakraborty and Mondal, 2021).

 $Log (COHb\%) = [0.858 \times log (CO)] + [0.63 \times log (t)] - 2.295$...(4)

2.5.5. Exposure assessment for NO₂ and SO₂

Equation Eq. 5 was used to determine the non-carcinogenic exposure (average hourly dose for inhalation $(\mu g/kg/hour)$) to NO₂ and SO₂ (Basu et al. 2024).

 $AHD = C \times IR/BW$

...(5)

For exposure to non-carcinogenic pollutants (NO₂ and SO₂), the average daily dose of the pollutant of interest ($\mu g/kg/day$) (Eq. 6) was calculated from the below formula:

	$ADD = (C \times IR \times ED)/(BW \times AT)$	(6)
6	Fransura assassment for NO. and SO.	

2.5.6. Exposure assessment for NO₂ and SO₂

The hazard quotient (HQ) was used to estimate the potential non-carcinogenic consequences of exposure to a known pollutant. The non-cancer likelihoods for cases of acute and chronic exposure were calculated using Eq. 7 and Eq. 8:

Chronic $HQ = ADD/REL$	(7)
Acute $HQ = AHD/REL$	(8)

Where REL (reference exposure level), as adopted from WHO (2021).

2.6. Statistical Analysis

The data were evaluated statistically utilising SPSS 27 and Microsoft Excel 13. The ANOVA was applied to analyse the difference between concentrations of pollutants from different fuels burning at a 5% significance level. The independent variable in this study is the socioeconomic condition of households, which encompasses factors such as caste, occupational structure, housing pattern, and income group. On the other hand, the dependent variable is the pattern of primary fuel usage. The association between socioeconomic status and cooking fuel use was investigated using the Chi-square (χ^2) test at 95 % significance level. To find out the uncertainty in health risk estimation, the Monte-Carlo simulation models were run for 10,000 simulations using Oracle Crystal Ball software version 11.

2.7. Analysis of land use/land cover

This study analysed the spatial distribution of land use in the studied area. To examine the correlation between the use of traditional fuels, such as firewood gathered from local forests, and socio-economic status, considering the availability of land use factors like vegetation, forests, and agricultural land. This objective aims to understand how access to traditional fuel sources is influenced by land use and how these sources, in turn, reflect and impact socio-economic conditions.

The present study involved the processing and analysis of a Sentinel-2A image acquired on 14th April 2023 at 11:54 a.m. UTC, as indicated in Supplementary Table 1a. The imagery was obtained from the Copernicus open access hub (https://scihub.copernicus.eu/), a platform that offers unrestricted open access to the user products of Sentinel-2 Level-1C (L_1C). L_1C products, alternatively referred to as tiles, encompass spatially referenced images spanning an area of 100 square kilometres. These images are projected in the Universal Transverse Mercator (UTM)/World Geodetic System 1984 projection-the tile designated as S2A_MSIL2A_20230414T043701_N0509_R033_T45QWG_20230414T081154.SAFE (Supplementary Table 1a) was selected for subsequent analysis due to its inclusion in our designated study site. Moreover, to identify the land use pattern in the study area, bands with 10-meter resolution have been selected, namely Band-2, Band-3, Band-4, and Band-8 (Supplementary Table 1b). Using a data management tool, all four bands have been composited in an Arc GIS environment (evaluation copy). The maximum likelihood classification method was applied to get the final output of the map. Five classes have been identified in the LULC map: water bodies, cultivated land, current fallow, built up and vegetation cover.

3. Results and Discussions

3.1. Socio-Economic Condition and Fuel Use

In India, rural households mainly depend on solid biomass fuel for their daily cooking needs, which is the source of HAP in the indoor environment, leading to severe health and environmental hazards. In contrast, biogas will be an alternative solution for climate change, and energy crises and to fulfil the increased energy demand due to population growth. Moreover, in recent decades, due to increased price hikes of LPG, which is a financial burden for rural households where LPG is used as a secondary energy option. The majority of the households in rural areas in India still rely on solid fuel. Several factors are associated with the choice of clean cooking fuel and only 0.4% of households used biogas as primary cooking fuel in India (Rajendran et al. 2012). In rural India, there are various types of biogas digesters, with most rural households using fixed dome biogas digesters. The Deenbandhu model, developed in 1984 by a voluntary organization, Action for Food Production, is a popular biogas digester among families due to its family-friendly design with various capacity options (Meena et al. 2024). Over time, KVIC has been actively promoting the Deenbandhu model as a biogas plant in rural areas of India, specifically at the community level (Singh et al., 2023). All the biogas plants in the study are fixed dome Deenbandhu model digesters with a capacity of 2 cubic meters. KVIC installed these digesters between 1994 and 1997. It is possible to generate one cubic meter of biogas per day, which is enough to meet the daily cooking energy needs of a family of 4-5 members (Aryal et al. 2022).

The socio-demographic characteristics of the village are provided in the Supplementary Table S1. The dominant economic status of the village (75%) households belongs to the lower-income group and lower-middle-income group. The households mainly engaged in daily wage earners account for almost 55.5%, and few households are engaged in the service sector (8%). The rest of the households engaged themselves in small businesses and other similar low-cost work. Hence, more than 66% of household heads were illiterate, and 76% of the household heads were male. In the village, most of the households are *pucca* (made up of brick and concreted roofs), and semi-*pucca* (made up of brick walls and asbestos of tin roof) comprise almost 43%, and the rest of the households are *Kuccha* structures made by mud and thatched roof (57%). Moreover, the village is covered by natural resources like trees, ponds, and agricultural land, but in the village, 38% of households are landless. The villagers belong to different social groups, i.e. General caste, Scheduled Caste (SC) and Scheduled Tribe (ST).

One of the key determinants of biogas as a primary fuel choice is the socio-economic conditions. In Table 1, chi-square statistics reveal the different cooking fuel types available based on the demographic characteristics of rural households. The present work discusses eight major socio-demographic characteristics that influence

the choice of cooking fuel in rural households, such as social category, household head, age, gender, education, housing patterns, occupational structure, and income groups. In the study area, 68.21% of households use firewood as their main cooking fuel source, and only a very small percentage (4.62%) of households use biogas as a source of cooking fuel, while others use cow dung cake and charcoal cake (27.17%). In contrast, LPG is used as an optional or secondary option with other energy sources. Most significantly, people were still using conventional fuel (95.38%), and only 4.62% of households used biogas as a primary fuel. There is a significant relationship between access to cooking fuel types and the social groups of rural households, as measured by the chi-square statistic ($\chi^2 = 31.594$, P < 0.01). The results indicate that households from general castes have the highest accessibility of biogas as a clean cooking fuel (17.95%), whereas only 1.52% of households access biogas; more interestingly, ST households don't use biogas as fuel. This is observed due to the caste inequality and lower economic status still considered a major hindrance to adopting biogas energy that prevails in rural Indian society (Bagdi et al. 2022; Soni and Chatterjee 2023). However, most households in the study areas are still structured as *Kuccha* households (57%), which has significant relationship between housing pattern and access to cooking fuel ($\chi^2 = 28.241$, P < 0.01) was observed with the majority of Kuccha households using firewood (79.80%), while Pucca households' people still using biogas as their primary cooking fuel. Similar observations were reported by Baqir et al. (2019), Afridi et al. (2023), and Kalli et al. (2022) and found that most of the *Kuccha* households primarily depend on traditional fuelwood for their daily energy requirement. Some studies indicate that household structure influences the adoption of biogas technology (Yasmin and Grundmann 2019). The household occupations are in the service sector (21.43%), and gardeners (52.94%) consume less firewood than those engaged in manual labour, driving, maid service, and small business. Households in the services category find it easier to afford cattle maintenance costs. However, livestock, particularly cattle, play an essential role in the livelihoods of rural populations, especially in developing countries. As a result, they often use cow dung cakes mixed with firewood as a substitute for cooking fuel. The findings indicate a strong association between cooking fuel types and the occupational structure of the village ($\gamma 2 = 39.260, P < 0.01$). The chi-square test results ($\chi 2 = 45.779$, P < 0.01) reveal a significant relationship between income and access to cooking fuel. It was demonstrated that higher-income people had greater access to biogas as a clean and sustainable energy source. This suggests that higher-income households have a greater possibility of getting access to clean cooking fuels like biogas than lower-income households. Multiple studies concluded that households with higher incomes are more likely to switch and adopt biogas as a greener energy source than households with lower incomes (Ghosh et al. 2020). It also demonstrates that the tendency of changing patterns towards biogas adoption is highly related to household income level, which decreases firewood use and shifts towards renewable energy resources or clean energy.

	Firewood	Cow dung cake	Charcoal cake	Biogas	Total	χ2
Occupational structure						39.260***
Daily wage earner	78.13	16.67	2.08	3.13	55.49	
Driver	84.62	15.38	0.00	0.00	7.51	
Gardener	52.94	23.53	23.23	0.00	9.83	
Maid	75.00	0.00	25.00	0.00	4.62	
Service	21.43	35.71	28.57	14.29	8.09	
Small business	56.00	16.00	16.00	12.00	14.45	
Social groups						31.594***
General caste	48.72	23.08	10.26	17.95	22.54	
Scheduled Caste (SC)	62.12	24.24	12.12	1.52	38.15	
Scheduled Tribe (ST)	85.29	8.82	5.88	0.00	39.31	
Housing pattern						28.241***
Kuccha	79.80	12.12	4.04	4.04	57.23	
Pucca	53.13	28.13	6.25	12.50	18.50	
Mixed	52.38	23.81	23.81	0.00	24.28	

Table 1: Cooking energy using pattern among rural households by socio-demographic characteristics

Income Groups

Lower Income Group	91.78	2.74	2.74	2.74	42.20	
Lower Middle-Income Group	55.93	30.51	10.17	3.39	34.10	
Upper Middle-Income Group	50.00	35.00	10.00	5.00	11.56	
Higher Income Group	38.10	19.05	28.57	14.29	12.14	
Education level						66.570***
Illiterate	81.74	13.91	3.48	0.87	66.47	
Primary level	56.67	26.67	13.33	3.33	17.34	
Secondary level	38.89	27.78	22.22	11.11	10.40	
Higher secondary level	0.00	20.00	40.00	40.00	5.78	
Household's Head Age						17.199***
Below 30	62.50	22.50	15.00	0.00	23.12	
30 years – below 60 years	73.86	19.32	4.55	2.27	50.87	
60 years and above	62.22	11.11	13.33	13.33	26.01	
Household's kitchen pattern						13. 5***
Kuchcha	74.02	16.54	7.87	1.57	26.58	
Pucca	52.17	21.74	13.04	13.04	73.42	
Household's agricultural land						14.222***
Agriculture landless	59.09	28.79	12.12	0.00	38.15	
Agriculture landowner	73.83	11.21	7.48	7.48	61.85	
Total households	68.21	17.92	9.25	4.62		
Below 30 02.30 22.30 13.00 0.00 23.12 30 years - below 60 years 73.86 19.32 4.55 2.27 50.87 60 years and above 62.22 11.11 13.33 13.33 26.01 Household's kitchen pattern 13. 5*** 1.57 26.58 Pucca 52.17 21.74 13.04 13.04 73.42 Household's agricultural land 28.79 12.12 0.00 38.15 Agriculture landless 59.09 28.79 12.12 0.00 38.15 Total households 68.21 17.92 9.25 4.62 20.00 Source: author's calculation from household survey (percentage) 9.25 4.62 20.00 28.70						

Significance level: *** p < 0.01, **p < 0.05,

Moreover, the level of education of the households is one of the significant factors for the adoption and access to clean fuel. The results also demonstrate a significant relationship between the education level of the Head of household and access to clean cooking energy ($\chi 2 = 66.570$, P < 0.01). This indicates that increasing education levels will significantly impact access to clean cooking energy. Although the analysis indicates a significant relationship between age in the household head and access to cooking fuels in rural households, as measured by the chi-square statistic ($\chi 2 = 17.199$, P < 0.01). It implies that the choice of clean cooking fuel increases with the household member's age due to concern about the family benefits of clean energy over traditional energy, where HAP is one of the major issues. The majority of the households have a *kuchcha* kitchen structure (73.42%), which is mainly made of bamboo and mud-made walls with roofs covered by rice straw, while *pucca* kitchens are made of bricks with concrete roofs, mainly inside the main building. The study found there is a strong association ($\chi 2 = 13.5$, P < 0.01) between kitchen structure with fuel using the pattern found biomass user households generally used the *kuchcha* pattern.

In contrast, biogas user groups are mainly used *pucca* structure. This might be due to their lower economic status and lack of proper knowledge about the effects of indoor air pollution. Finally, the result found a significant association between access to cooking fuel types and the availability of agricultural land, as measured by the chi-square statistic ($\chi 2 = 14.222$, P < 0.01). This implies households with agricultural land are interested in clean energy, particularly biogas, as they can use the bio-slurry as an organic fertilizer for their agricultural land and clean cooking fuel. The fertiliser that the biogas plant produces can replace 15-20% of chemical fertiliser with organic fertiliser, which will sustain agricultural productivity (Show et al. 2023; Karmakar et al. 2024; Koley et al. 2024a). Therefore, households with biogas plants profit more by utilising biogas as cooking energy and bio-slurry as fertiliser. Numerous studies proposed that a household biogas plant benefits from all aspects of household needs and is also environment-friendly (Sarker et al. 2020; Singh and Kalamdad 2021).

3.2. Indoor Air Quality and Health Risk Assessment

3.2.1. Status of indoor particulate matter (PM2.5) and carbon monoxide (CO) concentration

The concentrations of PM_{2.5} and CO are presented in Table 2. The inhalation of exceeded mass concentrations of particulate matter (PM) (Chakraborty and Mondal 2021) and gaseous pollutants (Basu et al. 2024) compared to WHO-prescribed indoor air quality standards are responsible for both chronic and acute health effects. The degree of exposure and health risk in rural women increases, particularly those who have used biomass fuel as a cooking fuel for a long duration of time due to inhalation of PM (PM₁₀, PM_{2.5}) and gaseous pollutants, a byproduct of biomass burning. Previous studies show that PM_{2.5} penetrate deep into the alveolar area while PM_{2.5-10} mainly accumulates in the tracheobronchial airways in the human respiratory system (Schroeter et al. 2006). Hypoxia affects organs with higher oxygen use, including the developing fetus (Hutter and Jaeggi 2010). In the present study, the PM_{2.5} concentration was observed highest for coal cake (293.53 ± 11.75 μ g/m³) followed by cow dung (205.83 ± 9.47 μ g/m³) > wood (185.06 ± 6.45 μ g/m³) > biogas (1.11 ± 0.16 μ g/m³). The ANOVA analysis also showed a significant (p < 0.001) difference in PM_{2.5} concentrations among different fuel user groups. The same trend in the output of ANOVA analysis was also observed for carbon monoxide (CO). However, the highest concentrations were monitored for cow dung users (6.46 ± 0.28 ppm), followed by coal cake (4.79 ± 0.21 ppm) > wood (4.48 ± 0.21 ppm) > biogas (0.01 ± 0.01 ppm) (Table 2). The result of the present study shows that biogas has significantly lower emissions for both PM_{2.5} and CO, which may be a positive aspect for efficient fuel use for rural and urban households.

Table 2: The concentration	of $PM_{2.5} (\mu g/m^3)$	and CO (ppm) from	different fuel burning
----------------------------	---------------------------	-------------------	------------------------

Pollutant	Wood	Cow Dung	Coal Cake	Biogas	F value	P value
PM _{2.5}	185.06 ±	$205.83 \pm$	$293.53 \pm$	1.11 ±	3363.59 (R-Sq = 99.45%)	< 0.001
	6.45	9.47	11.75	0.16		
CO	$4.48 \pm$	$6.46 \pm$	$4.79 \pm$	$0.01 \pm$	2682.43 (R-Sq = 99.31%)	< 0.001
	0.21	0.28	0.21	0.01		

3.2.2. Health risk assessment from indoor particulate matter $(PM_{2.5})$

Present study results of $PM_{2.5}$ exposure index and intake concentration (Table 3) were observed almost 170-274 times and 167- 268 times higher than the biogas users. The median exposure index calculated for wood, cow dung, coal cake, and biogas were 2.05, 2.26, 3.29 and 0.012, respectively. At the same time, median intake concentrations were 118.71, 132.2, 190.07 and 0.707, respectively. A higher level of $PM_{2.5}$ in traditional cookstoves is possibly due to inefficient biomass combustion in poorly designed traditional stoves. The PM emission from traditional biomass cookstoves is well-documented (Mitra et al. 2023). A meta-analysis was conducted by Atkinson et al. (2014), who highlighted that every 10 µg/m³ increase of $PM_{2.5}$ leads to a 1% increase in hospital admission. The hazardous quotient was calculated for $PM_{2.5}$ exposure via the inhalation route, and the results demonstrated that it exceeded greater than one for every traditional fuel except biogas. The representable HQ values were 7.91, 8.81, 12.67 and 0.047 for wood, cow dung, coal cake and biogas, respectively (Table 3). An elevated hazard quotient level for biomass users may be due to the excessive deposition of particles in the respiratory tract (Chalvatzaki et al. 2019). Thus, a higher intake of $PM_{2.5}$ can induce the early symptoms of cardiovascular diseases (Mitra et al. 2022).

Table 3: Exposure Index, Intake Concentration, and	Toxicological Risk from PM2.5 concen	tration among different fuel user groups
--	--------------------------------------	--

	TT 7 1	0 D	0 101	
	Wood	Cow Dung	Coal Cake	Biogas
Exposure Index (EI)				
Median	2.05	2.26	3.29	0.012
1st quartile	2.02	2.26	3.19	0.012
3rd quartile	2.08	2.33	3.36	0.013
Intake concentration (IC)				
Median	118.71	132.2	190.07	0.707
1st quartile	117.08	131.01	184.93	0.707
3rd quartile	120.36	134.81	194.14	0.771
Hazard Quotient (HQ)				
Median	7.91	8.81	12.67	0.047
1st quartile	7.81	8.73	12.33	0.047
3rd quartile	8.02	8.99	12.94	0.051

3.2.3. Status of indoor particulate matter (PM_{2.5}) and carbon monoxide (CO) concentration

Blood carboxyhemoglobin percentage (COHb%) may be used to estimate the amount of carbon monoxide present in a person's body. According to the Commission of the European Communities research in 1991, my-ocardial infarction (MI) could have been linked to greater levels of carbon monoxide inhalation. According to earlier clinical studies, blood COHb concentrations above the 3% threshold can cause carbon monoxide poisoning, which can eventually be lethal (ATSDR 2009). The blood COHb% was recorded highest for Cow dung cake users ($0.39 \pm 0.02\%$) followed by Coal cake ($0.28 \pm 0.02\%$) > Wood ($0.25 \pm 0.02\%$) and least for biogas users ($0.008 \pm 0.01\%$). The COHb% values showed a significant (p < 0.001) difference among the biogas and other fuel user groups (Table 4). Chakraborty and Mondal (2021) found comparable calculated blood COHb% varies from 0.2% to 0.85% in non-smokers and is elevated in smokers (10%) and heavy smokers (4%). Mitra et al. (2022) discovered comparable research findings and health consequences from CO when they evaluated the physical condition of women from the same region who were exposed to wood smoke.

Table 4: Percentage of blood Ca	arboxyhemoglobin (CO	OHb) of diffe	erent fuel user g	roups	
Stove categories	$Mean \pm SD$	t value	CI (95%)	<i>p</i> -value	
Wood	0.25 ± 0.02	41.35	0.23, 0.26	< 0.001	
Cow Dung Cake	0.39 ± 0.02	74.11	0.37, 0.39	< 0.001	
Coal Cake	0.28 ± 0.02	85.97	0.26, 0.28	< 0.001	
Biogas	0.008 ± 0.01				

3.2.4. Concentration of indoor Nitrogen dioxide (NO2) and Sulphur dioxide (SO2) and Health risk assessment

The concentrations of NO₂ emitted from different fuel burning were used to calculate HQ (Table 5). The HQ calculated for acute and chronic exposures showed no immediate and prolonged adverse health effects for women (HQ < 1.0). However, cooking with biomass fuels may show some long-term chronic health risks, as evidenced by our analysis. The highest mean NO₂ concentration was found for coal $(1.42 \pm 0.16 \,\mu\text{g/m}^3)$, followed by cow dung $(1.31 \pm 0.15 \,\mu\text{g/m}^3) > \text{wood} (1.20 \pm 0.12 \,\mu\text{g/m}^3) > \text{biogas} (1.13 \pm 0.04 \,\mu\text{g/m}^3)$. The NO₂ concentration showed a significant (p < 0.001) difference among the biogas and other fuel user groups except wood fuel. Present risk evaluation found that acute exposure to indoor NO₂ posed a minimal risk. Recent research, however, has indicated that modest amounts of NO₂ exposure can cause acute and obstructive lung illnesses. Some investigations also discovered a link between NO₂ levels and acute ischemic stroke. However, several research studies have found no statistically significant links between NO₂ exposure and human health (Andersen et al. 2012; Chen et al. 2012).

Table 5: Hazard quotients for acute and chronic exposure health risk from nitrogen dioxide (NO₂, μ g/m3) among different fuel user groups

Fuel types	$Mean \pm SD$	Acute HQ	Chronic HQ	<i>p</i> -value
Wood	1.20 ± 0.12	1.05×10 ⁻³	4.64×10 ⁻²	< 0.073
Cow dung	1.31 ± 0.15	1.15×10 ⁻³	5.09×10 ⁻²	< 0.001
Coal	1.42 ± 0.16	1.24×10 ⁻³	5.51×10 ⁻²	< 0.001
Biogas	1.13 ± 0.04	9.88×10 ⁻⁴	4.38×10 ⁻²	

This study is 1st of its kind, which seeks to explore the health risk assessment from SO₂ concentration from burning biomass and biogas fuels in indoor microenvironment. The concentrations of SO₂ emitted from different fuel burning were used to calculate HQ (Table 6). The HQ calculated for acute and chronic exposures showed no immediate and prolonged adverse health effects for women (HQ < 1.0). The highest means SO₂ concentration was found for coal ($0.055 \pm 0.008 \,\mu g/m^3$) followed by wood ($0.015 \pm 0.004 \,\mu g/m^3$) > cow dung ($0.008 \pm 0.002 \,\mu g/m^3$) > biogas ($0.002 \pm 0.001 \,\mu g/m^3$). The SO₂ concentration showed a significant (p < 0.001) difference

among the biogas and other fuel user groups. Chakraborty et al. (2014), as well as Mondal and Chakraborty (2015), also observed similar results. In contrast to biomass, many coals include inherent pollutants such as lead, mercury, silica, fluorine, sulfur, and arsenic. These pollutants are not eliminated by combustion; they are discharged into the atmosphere in their unaltered or oxidized state. For instance, sulfur dioxide (SO₂) pollution has a local or regional impact on outdoor and interior air quality in homes that burn sulfur-rich coal since coal burns at a significantly higher temperature than biomass combustion, which produces more NO₂ emissions than biomass combustion (Zhang et al. 2000).

1	1		1	, 0
	$Mean \pm SD$	Acute HQ	Chronic HQ	<i>p</i> -value
Wood	0.015 ± 0.004	8.20×10 ⁻⁶	3.63×10 ⁻⁴	< 0.001
Cow dung	0.008 ± 0.002	4.29×10 ⁻⁶	1.90×10 ⁻⁴	< 0.001
Coal	0.055 ± 0.008	3.01×10 ⁻⁵	1.33×10 ⁻³	< 0.001
Biogas	0.002 ± 0.001	1.02×10 ⁻⁶	4.51×10 ⁻⁵	

Table 6: Hazard quotients for acute and chronic exposure health risk from Sulphur dioxide (SO₂, μ g/m³) among different fuel user groups

3.2.4. Health Risk assessment and Monte-Carlo Probabilistic simulation

There is always a chance to have uncertainties in health risk estimation. The uncertainty may present among the concentrations of pollutants, variables used for risk estimation and/or other environmental factors. The median value of HQ for wood $PM_{2.5}$ was found 5.10 with a certainty range from 4.13 to 6.28 at a 95% level of confidence interval during the use of biomass cookstove (Fig. 2A). Whereas the median value of HQ for cow dung PM2.5 was observed at 5.67 with a certainty range from 4.53 to 7.04 at a 95% level of confidence interval in case of biomass users (Fig. 2B). However, the highest median value of HQ for coal cake $PM_{2.5}$ was found 8.08 with a certainty range from 6.54 to 10.00 at a 95% level of confidence interval during the use of traditional coal cookstove (Fig. 2C) and the median value of HQ for biogas $PM_{2.5}$ was found 0.03 with a certainty range from 0.02 to 0.04 at a 95% level of confidence interval during the use of confidence interval during the use of biogas server time and exposure duration were the two most important factors to represents in the health risk estimation for wood, cow dung and coal cake users, however, for biogas users $PM_{2.5}$ showed the highest contribution (Fig. 3A-3D). The above results may be due to high or low ventilation present in the particular households. The present study has observed that traditional cookstove users are more prone to non-carcinogenic health risks, whereas biogas represents a clean and safer cooking fuel.



Fig. 2: Monte-Carlo probabilistic distribution of the Hazard Quotient (HQ) for wood $PM_{2.5}$ (A), for cow dung $PM_{2.5}$ (B), for coal cake $PM_{2.5}$ (C) during the use of traditional biomass cookstove and for biogas $PM_{2.5}$ during the use of biogas cookstove (D)

The study's major contribution is the incorporating of self-reported health-related data related to household air pollution across various user groups, along with a comparative analysis of biogas user groups. Presently, due to energy shortages and environmental protection, biogas is an alternative solution to mitigate these upcoming threats. However, the study design has some limitations, such as the smaller sample size of biogas users compared to solid fuel user groups. The study only focused on a single village; it could be conducted in different parts of the location with regional variation. Thus, further study will provide more significant results. The study does not include other relevant socio-economic factors such as religion, livestock, gender, food habits, and household size. Future studies might carefully address other unmeasured pollutants like PAHs and benzene, along with other health-related parameters like BMI status, nutritional status, blood pressure, and heart rate.



Fig. 3: Sensitivity factors after running Monte-Carlo probabilistic distribution of the Hazard Quotient (HQ) for wood users (A), for cow dung users (B), for coal cake users (C), and for biogas (D) users.

3.2.5. Self-reported symptoms of health-related issues among different fuel user groups

Table 7 presents the self-reported health problems faced by households with different fuel user groups. Table 7 reveals that solid fuel biomass user groups primarily experience eye irritation, breathlessness, wheezing

cough, and muscle pain; conversely, only a small proportion of biogas users experience headache, sneezing, chest pain, and arthritis. The current study emphasized that muscle pain is a significant issue for fuelwood users, particularly women who handle loads of firewood exceeding 20 kg for extended periods. This issue affects 17% of women who use solid fuel, excluding those who use biogas.

According to the women's self-reported health problems, firewood users reported the highest percentage of headaches (50%) among the different fuel types. In comparison, biogas users had a significantly lower incidence (11.8%), with 18 individuals (10.4%) overall reporting headaches across all fuel types. Eye irritation was most common among firewood users (70%), with no cases reported among biogas users, indicating a strong association between traditional biomass fuels and eye irritation, as 20 individuals (11.6%) reported this symptom. Breathlessness was reported by 61.5% of firewood users and 26.9% of cow dung cake users, with no cases among biogas users, suggesting a health benefit of biogas use by 26 individuals (15%) who reported breathlessness. Cough was also most prevalent among firewood users (79.2%), with no cases among biogas users, contributing to the 24 individuals (13.9%) who reported this condition. Chest pain was reported by 61.1% of firewood users (77.4%) and absent among biogas users, with 31 individuals (17.9%) reporting this symptom. Sneezing was reported by 70.6% of firewood users (12 individuals) and less frequently by biogas users (5.6%), with a total of 17 cases (9.8%). Lastly, arthritis was reported by 68.4% of firewood users, contributing to the 19 cases (11%) overall.

Types	Firewood	Cow dung cake	Coal cake	Biogas	Total
Headache	50.0 (9)	16.7 (3)	22.2 (4)	11.8 (2)	10.4 (18)
Eye Irritation	70.0 (14)	20.0 (4)	10.0(2)	0.0 (0)	11.6 (20)
Breathlessness	61.5 (16)	26.9(7)	11.5 (3)	0.0 (0)	15 (26)
Cough	79.2 (19)	16.7 (4)	4.2 (1)	0.0 (0)	13.9 (24)
Chest pain	61.1 (11)	22.2 (4)	11.1 (2)	5.6(1)	10.4 (18)
Muscle pain	77.4 (24)	12.9 (4)	9.7 (3)	0.0 (0)	17.9 (31)
Sneezing	70.6 (12)	11.8 (2)	11.8 (2)	5.9(1)	9.8 (17)
Arthritis	68.4 (13)	15.8 (3)	10.5 (2)	5.3 (1)	11 (19)

Table 7: Classification of self-reported health status symptoms among different biomass user and biogas user groups

Matinga and Clancy (2020), as well as Koyuncu et al. (2021), observed that women were burdened with a significant burden of muscle pain and back pain. Women who use solid fuel primarily report eye irritation issues. This is possible due to the absence of proper ventilation and smoke generation from the burning of solid fuel biomass, which will sometimes enhance the risk of cataracts. A recent study by Kumar et al (2023) and Mandell et al (2020) highlighted that cataracts are common among solid fuel biomass user women in China. The issue doesn't impact biogas user groups because they burn biogas fuel without emitting smoke. Arthritis, also commonly known as joint pain symptoms, was highly preventable among solid fuel user groups. Liu et al. (2022) highlighted that exposure to PM_{2.5} increases the risk of arthritis among solid fuel biomass user groups. The questionnaire-based survey results revealed that solid fuel user groups experience more common respiratory issues such as whooping cough, breathlessness, and sneezing.

In contrast, only a small number of biogas user women reported experiencing sneezing problems. This observation aligns with a previous study which found that upper and lower respiratory systems affect 50.9% and 70.9% of premenopausal biomass-using women in India, respectively. Similarly, Wang et al. (2023) found a significant correlation between exposure to $PM_{2.5}$ and respiratory infection by the air pollutant. Similarly, Pathak et al. (2020) found a higher prevalence of respiratory symptoms among biomass user groups compared to clean fuel user groups. The study demonstrates a correlation between biomass exposure and an elevated risk

of respiratory system issues. Therefore, the current study indicates that households using solid fuels experience more health-related issues than those receiving biogas.

3.2.6. Health Risk and Socioeconomic condition

Socioeconomic conditions severely influence the health risks associated with traditional fuel use in rural areas. Traditional fuels like firewood, cow dung cake, and coal cake are prevalent in lower-income households due to their affordability and accessibility, but they pose significant health risks. These fuels emit high levels of particulate matter (PM_{2.5}) and carbon monoxide (CO), which are linked to severe respiratory and cardiovascular diseases. For instance, the exposure to PM_{2.5} in households using these fuels is drastically higher compared to those using biogas, leading to increased incidences of chronic health conditions.

The reliance on traditional fuels is often a result of socioeconomic factors such as income, education, and social status. In rural communities, where a large proportion of households fall into lower-income groups and depend on daily wage labour, the financial burden of cleaner fuels is often too high. Moreover, households from marginalized social groups, such as SC and ST, and those living in *kuccha* (mud) houses are more likely to use traditional biomass fuels. These groups often lack access to information about the health benefits of cleaner fuels and the economic resources needed to adopt them. Conversely, higher-income households, those with better education, and those living in pucca (brick) houses are more likely to use biogas, which significantly reduces health risks by emitting negligible amounts of harmful pollutants. However, despite the health and economic benefits of biogas, its adoption remains low in many rural areas due to persistent socioeconomic barriers.

3.2.7. Dependency on Traditional Fuels

In rural India, women still rely on biomass fuels like firewood, manure, and agricultural residues, even though biogas is a more sustainable and eco-friendly energy source. Rural women face many challenges when implementing biogas, including a reduced reliance on male household members for labour-intensive cooking fuel collection, which is often seen as an integral part of their domestic responsibilities. Cultural preferences, like as the taste of meals prepared over an open fire or firewood, are sometimes the cause of resistance to the adoption of biogas. Although biogas facilities need a large initial investment and continuous maintenance, financial restrictions are also a major factor. Furthermore, biomass is a reliable and readily stored energy source that ensures fuel security for rural women in areas where firewood is easily accessible.

In the study area, Sehalai village exhibits a dispersed settlement pattern (Fig. 4), with sample households covering an area of about 0.0494 km². The findings reveal evident spatial segregation of social groups, with the ST population located in a specific northern settlement. In contrast, SC and general caste households are situated in closer proximity to other groups. The village consists of 0.227 km² of waterbodies, which include ponds and canals, while a river flows through its southern area, providing sufficient water resources for agricultural activities. The area covered by homestead trees and woodlands is 0.6749 km², whereas the estimated cultivable land area is 2.0316 km². The presence of these natural resources contributes an essential role in shaping household energy decisions, as the accessibility of firewood supports the practice of fuel stacking. Women, who predominantly control cooking responsibilities, rely upon readily available biomass resources because alternative energy sources are often too expensive or scarce, thus continuing the reliance on traditional fuels.

In addition, the production of biogas frequently exhibits variability, especially during periods of limited feedstock supply or colder temperatures, causing households to shift back to biomass fuels to meet their energy requirements. Moreover, women residing in rural regions frequently encounter constraints in their decision-making control, especially when it comes to financial investments in clean cooking technologies. The transition to biogas is further hindered by a lack of awareness regarding the health benefits of cleaner fuels, alongside

insufficient understanding of the health risks linked to biomass combustion, which include cough, eye irritation, and respiratory illnesses.



Fig. 4: Land use Pattern of the North and South Sehalai village

4. POLICY RECOMMENDATION

To effectively involve the community in the implementation and upkeep of the biogas adoption, particularly in rural areas, by increasing awareness of the advantages of biogas, focusing on the effects on the environment and human health. To encourage government organisations, community-based organisations, non-governmental organisations, and other stakeholders to take effective actions to implement biogas-related schemes and policies.

Implementing biogas schemes in collaboration with income and employment-generating government programmes, such as Mahatma Gandhi National Rural Employment Guarantee Act, will have a more direct impact on the community. This will lead to reduced dependence on traditional cooking fuels. Even though there were subsidies for India's SC and ST population, they still had less biogas adoption due to a lack of subsidies, so the government focused more on subsidising marginalised and disadvantaged groups, especially the women members of economically and socially weaker sections of society.

5. CONCLUSION

The present study results revealed that rural households using traditional fuel showed a higher risk of health hazards directly and indirectly, particularly for women, due to the generation of household air pollution. Biogas is one of the feasible solutions to reduce HAP concentration, workload, health risks, and environmental threats by replacing traditional fuels. The present study found that biogas has emerged as a less polluting energy source (in terms of particulate matter and gaseous pollutants emission) than other traditional cooking energy sources. However, most of the population used it due to its easy availability and low cost. Moreover, factors such as income, age, education, occupation and social categories may play a significant role in choosing biogas as their

primary energy source. Moreover, the highest biomass smoke user groups observed increased muscle pain, eye irritation, cough and breathlessness when compared to biogas users. However, due to a lack of awareness about the technology used, rural people are still not ready to accept the biogas fuel, although it has great potential. Though the Government of India promote several schemes related to installing family-type biogas plants by MNRE in rural households, several drawbacks (e.g. technological barriers, fuel stacking, lack of supply chain and network system, unavailability of trained professional and awareness among SC and ST people) of the schemes which are required to take into consideration for policy formulation to achieve UN-SDG by 2030. The Government of India should facilitate proper initiatives to implement the schemes regarding the biogas plant at the household level to increase biogas adoption. The study mainly focuses on the women user group, while for future studies, a comparative study with health status in men among marginalized groups. The perceptional barriers to biogas adoption and certain health indicators like heart rate, blood pressure, and body mass index are not well covered in this study. Future studies should examine long-term advantages, include scientific evaluations of health effects, and concentrate on sociocultural elements impacting the use of biogas. Research on gender dynamics, the efficacy of policies, and supply chain constraints are required, as are comparisons with other clean energy sources. Furthermore, evaluating the advantages of biogas adoption on the environment and climate could help shape sustainable energy legislation.

Author Contributions: Tapas Bagdi, Apurba Koley, Deep Chakraborty: Conceptualization, Data curation, Investigation, Methodology, Software, Formal analysis, Writing - original draft; Sushil Kumar, Sreya Ghosh and Anuradha Sarkar: Data curation, Investigation; Writing - original draft, review; Niladri Das: Visualisation, Software and analysis; Suman Das and Sujan Gayen: Data collection and curation, Resources; Amit Kumar Hazra, Shibani Chaudhury, Andrew B Ross, Srinivasan Balachandran: Funding acquisition, Conceptualization, Supervision, Writing - review and editing.

Funding: This work was supported by BBSRC, United Kingdom, for receiving funding from the BEFWAM project: Bioenergy, Fertilizer and Clean water from Invasive Aquatic macrophytes [Grant Ref: BB/S011439/1]

Institutional Review Board Statement: For this study, ethical approval (VB/IECHR/07/2024) was granted by the Institutional Ethics Committee for Human Research of Visva-Bharati (A Central University), West Bengal, India.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Acknowledgements: Tapas Bagdi, Apurba Koley and Sreya Ghosh are thankful to the BBSRC, United Kingdom, for receiving funding from the BEFWAM project: Bioenergy, Fertilizer and Clean water from Invasive Aquatic macrophytes [Grant Ref: BB/S011439/1] for financial support and research fellowship.

Conflicts of Interest: The authors declare no conflicts of interest.

REFERENCES

- Acheampong, A. O., Opoku, E. E. O., and Dogah, K. E., 2023. The Political Economy of Energy Transition: The Role of Globalization and Governance in the Adoption of Clean Cooking Fuels and Technologies. *Technological Forecasting and Social Change*, 186, 122156. <u>https://doi.org/10.1016/j.techfore.2022.122156</u>.
- Afridi, F., Debnath, S., Dinkelman, T., and Sareen, K., 2023. Time for Clean Energy? Cleaner Fuels and Women's Time in Home Production. *The World Bank Economic Review*, 37(2), pp.283–304. <u>https://doi.org/10.1093/wber/lhac031</u>.
- Ahmad, M., and Wu, Y., 2022. Household-Based Factors Affecting Uptake of Biogas Plants in Bangladesh: Implications for Sustainable Development. *Renewable Energy*, 194, pp. 858–867. https://doi.org/10.1016/j.renene.2022.05.135.
- Akter, S., Mathew, N. M., and Fila, M. E., 2023. The impact of an improvement in the quality and reliability of rural residential electricity supply on clean cooking fuel adoption: Evidence from six energy poor Indian states. *World Development*, 172, 106366. <u>https://doi.org/10.1016/j.worlddev.2023.106366</u>.

- Alexander, S., and Padmanabhan, V., 2019. With Ujjwala, more LPG access but limited usage. *The Mint.* Retrieved June 3, 2024 from <u>https://www.livemint.com/news/india/with-ujjwala-more-lpg-access-but-limited-usage-1553571493940.html.
 </u>
- Ali, A., Mottaleb, K. A., and Aryal, J. P., 2019. Wealth, Education and Cooking-Fuel Choices among Rural Households in Pakistan. *Energy Strategy Reviews*, 24, pp.236–243. <u>https://doi.org/10.1016/j.esr.2019.03.005</u>.
- Ali, J., and Khan, W., 2022. Factors affecting access to clean cooking fuel among rural households in India during COVID-19 pandemic. *Energy for Sustainable Development*, 67, pp.102–111. https://doi.org/10.1016/j.esd.2022.01.006.
- Awan, A., Bilgili, F., and Rahut, D. B., 2023. Household fuel choices and consumption intensity in Pakistan: Evidence from HIES data 2001–2019. *Environmental Science and Pollution Research*, pp.1-16. https://doi.org/10.1007/s11356-023-27227-3.
- 9. Aziz, S., Barua, S., and Chowdhury, S. A., 2022. Cooking Energy Use in Bangladesh: Evidence from Technology and Fuel Choice. *Energy*, 250, 123696. <u>https://doi.org/10.1016/j.energy.2022.123696</u>.
- Bagdi, T., Ghosh, S., Sarkar, A., Hazra, A. K., Balachandran, S., and Chaudhury, S., 2022. Whose development counts? Adoption of biogas in the rural communities of India-a review. *International Journal of Renewable Energy Research*, 12(4), pp.2023-2042. <u>https://doi.org/10.20508/ijrer.v12i4.13451.g8561</u>.
- Bagdi, T., Ghosh, S., Sarkar, A., Hazra, A. K., Balachandran, S., & Chaudhury, S. (2023). Evaluation of research progress and trends on gender and renewable energy: A bibliometric analysis. *Journal of Cleaner Production*, 138654. <u>https://doi.org/10.1016/j.jclepro.2023.138654</u>
- Bai, C., Zhan, J., Wang, H., Yang, Z., Liu, H., Liu, W., Wang, C., and Chu, X., 2023. Heating choices and residential willingness to pay for clean heating: Evidence from a household survey in rural China. *Energy Policy*, 178, 113617. https://doi.org/10.1016/j.enpol.2023.113617.
- Balakrishnan, K., Dey, S., Gupta, T., Dhaliwal, R. S., Brauer, M., Cohen, A. J., Stanaway, J. D., Beig, G., Joshi, T. K., Aggarwal, A. N., et al., 2019. The impact of air pollution on deaths, disease burden, and life expectancy across the states of India: The Global Burden of Disease Study 2017. *The Lancet Planetary Health*, 3(1), pp.26-39. https://doi.org/10.1016/S2542-5196(18)30261-4.
- Baqir, M., Bharti, S. K., Kothari, R., and Singh, R. P., 2019. Assessment of an Energy-Efficient Metal Chulha for Solid Biomass Fuel and Evaluation of Its Performance. *International Journal of Environmental Science and Technol*ogy, 16(11), pp.6773–6784. <u>https://doi.org/10.1007/s13762-018-2028-9</u>.
- 15. Biswas, S., and Das, U., 2022. Adding Fuel to Human Capital: Exploring the Educational Effects of Cooking Fuel Choice from Rural India. *Energy Economics*, 105, 105744. <u>https://doi.org/10.1016/j.eneco.2021.105744</u>.
- Carley, S., and Konisky, D. M., 2020. The justice and equity implications of the clean energy transition. *Nature Energy*, 5(8), pp.569-577. <u>https://doi.org/10.1038/s41560-020-0641-6</u>.
- 17. CEDA (2023). How many Indians live in a pucca house with basic amenities? Retrieve on March 12, 2024 from https://ceda.ashoka.edu.in/how-many-indians-live-in-a-pucca-house-with-basic-amenities/
- Chanchani, D., and Oskarsson, P., 2021. If the gas runs out, we are not going to sleep hungry: Exploring household energy choices in India's critically polluted coal belt. *Energy Research and Social Science*, 80, 102181. https://doi.org/10.1016/j.erss.2021.102181.
- Chand, S., and Behera, B., 2023. Does assignment of individual property rights improve forest conservation outcomes? Empirical evidence from West Bengal, India. *Ecology, Economy and Society–the INSEE Journal*, 6(2), pp.7-31. <u>https://doi.org/10.37773/ees.v6i2.709</u>.
- Chen, T., Zou, C., Yuan, Y., Pan, J., Zhang, B., Qiao, L., Li, Y., Qian, J., Guo, Q., Yuan, Y., and Ding, C. 2022. Indoor air pollution from solid fuel on children pneumonia in low-and middle-income countries: A systematic review and meta-analysis. *Environmental Science and Pollution Research International*, pp.1-15. <u>https://doi.org/10.1007/s11356-021-18293-6</u>.
- 21. Chicombo, A. F. F., and Musango, J. K., 2022. Towards a theoretical framework for gendered energy transition at the urban household level: A case of Mozambique. *Renewable and Sustainable Energy Reviews*, 157, 112029. https://doi.org/10.1016/j.rser.2021.112029.
- Chowdhury, M., Ghosh, S., and Padhy, P. K., 2022. Effects of indoor air pollution on tribal community in rural India and health risk assessment due to domestic biomass burning: A realistic approach using the lung deposition model. *Environmental Science and Pollution Research International*, pp.1-13. <u>https://doi.org/10.1007/s11356-022-19973-7</u>.
- Ciplet, D. 2021. From energy privilege to energy justice: A framework for embedded sustainable development. *Energy Research and Social Science*, 75, 101996. <u>https://doi.org/10.1016/j.erss.2021.101996</u>.

- 24. Council on Energy, Environment and Water (CEEW). 2020. State of clean cooking energy access in India. Retrieved from https://www.ceew.in/publications/state-of-clean-cooking-energy-access-in-india-ires-2020-report.
- Council on Energy, Environment and Water (CEEW). 2021. More than 70% of Indian households use LPG as primary cooking fuel. Retrieved from <u>https://www.ceew.in/press-releases/more-70-indian-households-use-lpg-primary-cooking-fuel-ceew</u>.
- Das, S., Groote, H. D., and Behera, B., 2014. Determinants of Household Energy Use in Bhutan. *Energy*, 69, pp.661–672. <u>https://doi.org/10.1016/j.energy.2014.03.062</u>.
- Dida, G. O., Lutta, P. O., Abuom, P. O., Mestrovic, T., and Anyona, D. N., 2022. Factors predisposing women and children to indoor air pollution in rural villages, Western Kenya. *Archives of Public Health*, 80(1), pp.1-13. https://doi.org/10.1186/s13690-022-00791-9.
- 28. Faizan, M. A., and Thakur, R., 2019. Association between solid cooking fuels and respiratory disease across sociodemographic groups in India. *Journal of Health Pollution*, 9(23).
- Fanari, E. 2022. Struggles for just conservation: An analysis of India's biodiversity conservation conflicts. *Journal of Political Ecology*, 28(1). <u>https://journals.librarypublishing.arizona.edu/jpe/article/5214/galley/5096/download/</u>.
- 30. Foysal, M. A., Hossain, M. L., Sultana, A. R. S., Uddin, M. K., Sayem, M. M., and Akhter, J., 2012. Household energy consumption pattern in rural areas of Bangladesh. *Indian Journal of Energy*, 1(5), 72-85.
- Fuller, S., and Bouzarovski, S., 2023. Introduction to the Handbook on Energy Justice. In Handbook on Energy Justice Edward Elgar Publishing. Pp.1–12. Retrieve from <u>https://books.google.com/books?hl=en&lr=&id=tki-EAAAQBAJ&oi=fnd&pg=PA1&dq=Three+fundamental+components+of+energy+justice+are+distributional,+procedural,+and+recognition+&ots= L29jn 5Ex&sig=SBJsFhxlLLFLj staYS5QqiU7gE.
 </u>
- Gould, C. F., and Urpelaine, J., 2018. LPG as a clean cooking fuel: Adoption, use, and impact in rural India. *Energy Policy*, 122, pp.395-408. <u>https://doi.org/10.1016/j.enpol.2018.07.042</u>.
- Gould, C. F., Hou, X., Richmond, J., Sharma, A., and Urpelainen, J., 2020. Jointly modeling the adoption and use of clean cooking fuels in rural India. *Environmental Research Communications*, 2(8). <u>https://doi.org/10.1088/2515-7620/abaca9</u>.
- Gould, C. F., Schlesinger, S. B., Molina, E., Bejarano, M. L., Valarezo, A., and Jack, D. W., 2020. Household fuel mixes in peri-urban and rural Ecuador: Explaining the context of LPG, patterns of continued firewood use, and the challenges of induction cooking. *Energy Policy*, 136, 111053. <u>https://doi.org/10.1016/j.enpol.2019.111053</u>.
- Gupta, D., Sinha, M., and Chhatre, A., 2022. India's Forest Rights Act and Indigenous claims to community forest resources: A case study of Lavari, Maharashtra. World Development Perspectives, 27, 100449. <u>https://doi.org/10.1016/j.wdp.2022.100449</u>.
- Habib, G., Kumari, J., Khan, M., Imran, M., Zaidi, K., Yogesh, A., Nagendra, S., et al. 2023. Estimating shifts in fuel stacking among solid biomass fuels and liquified petroleum gas in rural households: A pan-India analysis. *Research Square*. <u>https://doi.org/10.21203/rs.3.rs-2674609/v1</u>.
- 37. International Energy Agency (IEA). 2021. Energy in India today India energy outlook 2021 analysis. Retrieved from https://www.iea.org/reports/india-energy-outlook-2021/energy-in-india-today.
- Irwin, B. R., Hoxha, K., and Grépin, K. A., 2020. Conceptualising the effect of access to electricity on health in lowand middle-income countries: A systematic review. *Global Public Health*, 15(3), pp.452-473.
- 39. Jenkins, K., McCauley, D., Heffron, R., Stephan, H., and Rehner, R., 2016. Energy justice: A conceptual review. *Energy Research and Social Science*, 11, pp.174-182. <u>https://doi.org/10.1016/j.erss.2015.10.004</u>.
- Joon, V., Chandra, A., and Bhattacharya, M., 2009. Household energy consumption pattern and sociocultural dimensions associated with it: A case study of rural Haryana, India. *Biomass and Bioenergy*, 33(11), pp.1509-1512. https://doi.org/10.1016/j.biombioe.2009.07.016.
- Karmakar, S., Koley, A., Thakur, R. G., Ghosh, A., Show, B. K., & Balachandran, S. 2024. Optimizing Hydroponic Growth by the Liquid Fraction of Biogas Digestate. In Changing Scenario of Earth, RAJESH PUBLICATIONS, New Delhi - 110002; ISBN: 978-93-91139-53-7
- 42. Kalli, R., Jena, P. R., and Managi, S., 2022. Subsidized LPG scheme and the shift to cleaner household energy use: Evidence from a tribal community of Eastern India. *Sustainability*, 14(4), 2450. <u>https://doi.org/10.3390/su14042450</u>.
- Kelkar, G., and Nathan, D., 2021. Cultural and economic barriers in switching to clean cooking energy: Does women's agency make a difference? *Energies*, 14(21), 7242. <u>https://doi.org/10.3390/en14217242</u>.
- Khandker, S. R., Barnes, D. F., and Samad, H. A., 2012. Are the energy poor also income poor? Evidence from India. *Energy Policy*, 47, pp.1-12. <u>https://doi.org/10.1016/j.enpol.2012.02.028</u>.

- 45. Khurana, T., and Sangita, S., 2020. Household access to electricity and non-farm business in rural India: A panel data analysis. *Energy for Sustainable Development*, 67, 125-134. <u>https://doi.org/10.1016/j.esd.2022.01.008</u>.
- 46. Koley, A.; Mukhopadhyay, P.; Show, B.K.; Ghosh, A.; Balachandran, S.; Santiniketan, V.-B. Biogas Production Potentiality of Water Hyacinth, Pistia and Duckweed: A Comparative Analysis. In Proceedings of the National Symposium: "Recent Trends in Sustainable Technology-Techno-Commercial Developments", Kolkata, India, 9–10 September 2022
- Koley, A., Bray, D., Banerjee, S., Sarkar, S., Thahur, R. G., Hazra, A. K., ... & Balachandran, S. (2022b). Water hyacinth (Eichhornia crassipes) a sustainable strategy for heavy metals removal from contaminated waterbodies. In *Bioremediation of Toxic Metal (loid) s* (pp. 95-114). CRC Press. <u>https://doi.org/10.1201/9781003229940-6</u>
- Koley, A., Mukhopadhyay, P., Gupta, N., Singh, A., Ghosh, A., Show, B. K., ... & Balachandran, S. (2023). Biogas production potential of aquatic weeds as the next-generation feedstock for bioenergy production: a review. *Environmental Science and Pollution Research*, 30(52), 111802-111832. <u>https://doi.org/10.1007/s11356-023-30191-7</u>
- Koley, A., GhoshThakur, R., Das, K., Gupta, N., Banerjee, A., Show, B. K., ... & Balachandran, S. (2024a). Growth Dynamics and Nutrient Removal from Biogas Slurry Using Water Hyacinth. Sustainability, 16(11), 4450. <u>https://doi.org/10.3390/su16114450</u>
- Koley, A., Gupta, N., Singh, A., GhoshThakur, R., & Balachandran, S. 2024b. Microalgae-Based Biofuel for Sustainable Bioenergy Production. In *Application of Microbial Technology in Wastewater Treatment and Bioenergy Recovery* (pp. 359-384). Singapore: Springer Nature Singapore. <u>https://doi.org/10.1007/978-981-97-3458-0-14</u>
- Nahar, G., Koley, A., Garai, S., Balachandran, S., & Ross, A. B. (2024). Enhancing Biogas Production Through the Co-Digestion of Fish Waste (FW) and Water Hyacinth (WH) Using Cow Dung as an Inoculum: Effect of FW/WH Ratio. *Applied Sciences*, 14(21), 9880. <u>https://doi.org/10.3390/app14219880</u>
- Kopas, J., York, E., Jin, X., Harish, S. P., Kennedy, R., Shen, S. V., and Urpelainen, J., 2020. Environmental justice in India: Incidence of air pollution from coal-fired power plants. *Ecological Economics*, 176, 106711. <u>https://doi.org/10.1016/j.ecolecon.2020.106711</u>.
- Kumar, K., and Sen, A., 2021. Socioeconomic development in highway corridor zones of Bihar: Case study of Kuchai Kot-Muzaffarpur section. In Resource Management, Sustainable Development and Governance. pp.563-579. <u>https://doi.org/10.1007/978-3-030-85839-1_35</u>.
- Li, N., Song, Q., Su, W., Guo, X., Wang, H., Liang, Q., Liang, M., Qu, G., Ding, X., Zhou, X., and Sun, Y., 2022. Exposure to indoor air pollution from solid fuel and its effect on depression: A systematic review and meta-analysis. *Environmental Science and Pollution Research International*, pp.1-15. <u>https://doi.org/10.1007/s11356-022-20841-7</u>.
- Ma, C., and Liao, H., 2018. Income elasticity of cooking fuel substitution in rural China: Evidence from population census data. *Journal of Cleaner Production*, 199, pp.1083–1091. <u>https://doi.org/10.1016/j.jclepro.2018.06.215</u>.
- Ma, W., Vatsa, P., and Zheng, H., 2022. Cooking Fuel Choices and Subjective Well-Being in Rural China: Implications for a Complete Energy Transition. *Energy Policy*, 165, 112992. <u>https://doi.org/10.1016/j.enpol.2022.112992</u>.
- 57. Majhi, S. 2022. A comparative analysis of availability of amenities between Scheduled Caste household and Non-Scheduled Caste household in West Bengal. *SHINJAN*, 21(8).
- Majumdar, D., Koley, M., and Chatterjee, P., 2023. Socioeconomic Factors and Clean Cooking Fuel Consumption in India: A Household Level Study. *Energy for Sustainable Development*, 76, 101298. <u>https://doi.org/10.1016/j.esd.2023.101298</u>.
- Malakar, Y., Greig, C., and Fliert, E. V., 2018. Resistance in rejecting solid fuels: Beyond availability and adoption in the structural dominations of cooking practices in rural India. *Energy Research and Social Science*, 46, pp.225-235. <u>https://doi.org/10.1016/j.erss.2018.07.025</u>.
- Meried, E. W. 2021. Rural Household Preferences in Transition from Traditional to Renewable Energy Sources: The Applicability of the Energy Ladder Hypothesis in North Gondar Zone. *Heliyon*, 7(11). <u>https://www.cell.com/heli-yon/pdf/S2405-8440(21)02521-4.pdf</u>.
- 61. Mosse, D. 2018. Caste and development: Contemporary perspectives on a structure of discrimination and advantage. *World Development*, 110, pp.422-436. <u>https://doi.org/10.1016/j.worlddev.2018.06.003</u>.

- 62. Mottaleb, K. A., Rahut, D. B., Aryal, J. P., and Ali, A., 2022. Clean fuel for rural families in India a major challenge: Evidence from four rounds of consumer expenditure survey. *Energy Reports*, 8, pp.2530–2546. https://doi.org/10.1016/j.egyr.2022.01.136.
- 63. Ngarava, S., Zhou, L., Ningi, T., Chari, M. M., and Mdiya, L., 2022. Gender and Ethnic Disparities in Energy Poverty: The Case of South Africa. *Energy Policy*, 161, 112755. <u>https://doi.org/10.1016/j.enpol.2021.112755</u>.
- Pachauri, S. 2007. An analysis of cross-sectional variations in total household energy requirements in India using micro survey data. *Energy Policy*, 32(15), pp.1723-1735. <u>https://doi.org/10.1016/j.jclepro.2018.12.092</u>.
- Pachauri, S. 2024. An analysis of cross-sectional variations in total household energy requirements in India using micro survey data. *Energy Policy*, 32(15), pp.1723-1735. <u>https://doi.org/10.1016/S0301-4215(03)00162-9</u>.
- Pachauri, S., and Jiang, L., 2008. The household energy transition in India and China. *Energy Policy*, 36(11), pp.4022-4035. <u>https://doi.org/10.1016/j.enpol.2008.06.016</u>.
- Patil, R., Roy, S., Gore, M., Ghorpade, M., Pillarisetti, A., Chakma, J., and Juvekar, S., 2021. Barriers to and facilitators of uptake and sustained use of LPG through the PMUY in tribal communities of Pune district. *Energy for Sustainable Development*, 63, pp.1–6. <u>https://doi.org/10.1016/j.esd.2021.04.008</u>.
- Patnaik, S., and Jha, S., 2020. Caste, class and gender in determining access to energy: A critical review of LPG adoption in India. *Energy Research and Social Science*, 67, 101530. <u>https://doi.org/10.1016/j.erss.2020.101530</u>.
- 69. Pelz, S., Chindarkar, N., and Urpelainen, J., 2021. Energy Access for Marginalized Communities: Evidence from Rural North India, 2015–2018. *World Development*, 137, 105204. <u>https://doi.org/10.1016/j.worlddev.2020.105204</u>.
- Perros, T., Allison, A. L., Tomei, J., and Parikh, P., 2022. Behavioural factors that drive stacking with traditional cooking fuels using the COM-B model. *Nature Energy*, 7(9), pp.886–898. <u>https://doi.org/10.1038/s41560-022-01074-</u>
 <u>x</u>.
- Petrokofsky, G., Harvey, W. J., Petrokofsky, L., and Ochieng, C. A., 2021. The importance of time-saving as a factor in transitioning from woodfuel to modern cooking energy services: A systematic map. *Forests*, 12(9). <u>https://doi.org/10.3390/f12091149</u>.
- Radhakrishnan, V., and Nihalani, J., 2022. Data | LPG prices skyrocket to ₹999.5, cylinder usage in rural areas under 45%. *The Hindu*. Retrieve on June 21, 2024 from <u>https://www.thehindu.com/data/data-lpg-prices-skyrocket-to-9995-cylinder-usage-in-rural-areas-under-45/article65398337.ece</u>.
- Radhakrishnan, V., Sen, S., and Nihalani, J., 2021. Most homes have LPG connection, but a significant share don't use it. *The Hindu*. Retrieved on June 21, 2024 from <u>https://www.thehindu.com/data/data-most-homes-have-lpg-connection-but-a-significant-share-dont-use-it/article34855341.ece</u>.
- Raghavendra, R. H., 2020. Literacy and health status of Scheduled Castes in India. *Contemporary Voice of Dalit*, 12(1), pp.97-110. <u>https://doi.org/10.1177/2455328X19898449</u>.
- Ranjan, R., and Singh, S., 2020. Household Cooking Fuel Patterns in Rural India: Pre- and Post-Pradhan Mantri Ujjwala Yojana. *Indian Journal of Human Development*, 14(3), pp.518–526. https://doi.org/10.1177/0973703020975045.
- Rao, M. N., and Reddy, B. S., 2007. Variations in energy use by Indian households: An analysis of micro level data. *Energy*, 32(2), pp.143-153. <u>https://doi.org/10.1016/j.energy.2006.03.012</u>.
- Ravindra, K., Kaur-Sidhu, M., Mor, S., and John, S., 2019. Trend in household energy consumption pattern in India: A case study on the influence of sociocultural factors for the choice of clean fuel use. *Journal of Cleaner Production*, 213, pp.1024-1034. <u>https://doi.org/10.1016/j.jclepro.2018.12.092</u>.
- Saxena, V., and Bhattacharya, P. C., 2018. Inequalities in LPG and electricity consumption in India: The role of caste, tribe, and religion. *Energy for Sustainable Development*, 4, pp.44-53. <u>https://doi.org/10.1016/j.esd.2017.09.009</u>.
- Setyowati, A. B. 2021. Mitigating inequality with emissions? Exploring energy justice and financing transitions to low carbon energy in Indonesia. *Energy Research and Social Science*, 71, 101817. <u>https://doi.org/10.1016/j.erss.2020.101817</u>.
- Sharma, D., Ravindra, K., Kaur, M., Prinja, S., and Mor, S., 2020. Cost evaluation of different household fuels and identification of the barriers for the choice of clean cooking fuels in India. *Sustainable Cities and Society*, 52, 101825. <u>https://doi.org/10.1016/j.scs.2019.101825</u>.
- Sharma, V., and Dash, M., 2022. Delineating energy consumption behaviour: A household-level assessment from India's Energy NEXUS Strategy. *Energy Nexus*, 100085. <u>https://doi.org/10.1016/j.nexus.2022.100085</u>.
- Show BK, Panja S, Ghosh Thakur R, Basu A, Koley A, Ghosh A et al (2023) Optimisation of anaerobic digestate and chemical fertiliser application to enhance rice yield—a machine-learning approach. *Sustainability* 15:13706 <u>https://doi.org/10.3390/su151813706</u>

- 83. Shrestha, R. P., Jirakiattikul, S., Lohani, S. P., and Shrestha, M., 2023. Perceived impact of electricity on productive end use and its reality: Transition from electricity to income for rural Nepalese women. *SSRN*, 4372924.
- Singh, K., Meena, R. S., Kumar, S., Dhyani, S., Sheoran, S., Singh, H. M., Pathak, V. V., et al. 2023. India's renewable energy research and policies to phase down coal: Success after Paris Agreement and possibilities post-Glasgow Climate Pact. *Biomass and Bioenergy*, 177, 106944. <u>https://doi.org/10.1016/j.biombioe.2023.106944</u>.
- 85. Somanaboina, S. 2021. The Other Backward Classes Pre-and Post-Mandal India. In The Routledge Handbook of the Other Backward Classes in India: Thought, Movements and Development.
- Soni, A., and Chatterjee, A., 2023. Not Just Income: The Enabling Role of Institutional Confidence and Social Capital in Household Energy Transitions in India. *Energy Research and Social Science*, 8, 103020. https://doi.org/10.1016/j.erss.2023.103020.
- Sovacool, B. K., and Dworkin, M. H., 2015. Energy justice: Conceptual insights and practical applications. Applied *Energy*, 142, pp.435-444. <u>https://doi.org/10.1016/j.apenergy.2015.01.002</u>.
- 88. Sovacool, B. K., Heffron, R. J., McCauley, D., and Goldthau, A., 2016. Energy decisions reframed as justice and ethical concerns. *Nature Energy*, 1(5), pp.1-6. <u>https://doi.org/10.1038/nenergy.2016.24</u>.
- Stoner, O., Lewis, J., Martínez, I. L., Gumy, S., Economou, T., and Adair-Rohani, H., 2021. Household cooking fuel estimates at global and country level for 1990 to 2030. *Nature Communications*, 12(1). <u>https://doi.org/10.1038/s41467-021-26036-x</u>.
- Swain, S. S., and Mishra, P., 2021. How does cleaner energy transition influence standard of living and natural resources conservation? A study of households' perceptions in rural Odisha, India. *Energy*, 215. https://doi.org/10.1016/j.energy.2020.119135.
- 91. USEPA 1989. Risk assessment guidance for superfund I human health evaluation manual (Part A) office of emergency and remedial response, Washington, DC.
- 92. USEPA 2005. Integrated risk information system. Adherence to long term therapies: evidence for action. https://www.epa.gov/iris.
- WHO. 2022. New global data on the use of clean and polluting fuels for cooking by fuel type. Retrieved January 2, 2024, from <u>https://www.who.int/news/item/20-01-2022-who-publishes-new-global-data-on-the-use-of-clean-and-polluting-fuels-for-cooking-by-fuel-type</u>.
- Williams, K. N., Kephart, J. L., Fandiño-Del-Rio, M., Nicolaou, L., Koehler, K., Harvey, S. A., and Checkley, W., 2023. Sustained use of liquefied petroleum gas following one year of free fuel and behavioral support in Puno, Peru. *Energy for Sustainable Development*, 73, pp.13–22. <u>https://doi.org/10.1016/j.esd.2023.01.005</u>.
- Yawale, S. K., Hanaoka, T., and Kapshe, M., 2021. Development of Energy Balance Table for Rural and Urban Households and Evaluation of Energy Consumption in Indian States. *Renewable and Sustainable Energy Reviews*, 136, 110392. <u>https://doi.org/10.1016/j.rser.2020.110392</u>.